

Rec'd PGT/PTO 18 APR 2005

**ALPHA MACROGLOBULIN FAMILY MEMBERS**

This invention relates to novel proteins, termed INSP087 and INSP088, herein identified as members of the alpha macroglobulin family, in particular as alpha-2-macroglobulin-like proteinase inhibitors, and to the use of these proteins and nucleic acid sequences from the  
5 encoding genes in the diagnosis, prevention and treatment of disease.

All publications, patents and patent applications cited herein are incorporated in full by reference.

**BACKGROUND**

The process of drug discovery is presently undergoing a fundamental revolution as the era  
10 of functional genomics comes of age. The term "functional genomics" applies to an approach utilising bioinformatics tools to ascribe function to protein sequences of interest. Such tools are becoming increasingly necessary as the speed of generation of sequence data is rapidly outpacing the ability of research laboratories to assign functions to these protein sequences.

15 As bioinformatics tools increase in potency and in accuracy, these tools are rapidly replacing the conventional techniques of biochemical characterisation. Indeed, the advanced bioinformatics tools used in identifying the present invention are now capable of outputting results in which a high degree of confidence can be placed.

Various institutions and commercial organisations are examining sequence data as they  
20 become available and significant discoveries are being made on an on-going basis. However, there remains a continuing need to identify and characterise further genes and the polypeptides that they encode, as targets for research and for drug discovery.

**ALPHA MACROGLOBULIN FAMILY**

The alpha macroglobulin family of proteins is divided into two general divisions - the  
25 alpha-2-macroglobulin like proteins and the complement-like proteins- that are thought to have arisen from a common ancestral alpha-2-macroglobulin-like molecule (Lin *et al*, 2002). The alpha macroglobulin family is therefore also known as the  $\alpha$ 2M/C3,C4,C5 family of thioester-containing protease inhibitor and complement proteins. A new member of the macroglobulin family, CD109, has recently been identified which has not yet been  
30 characterised as belonging to either the alpha-2-macroglobulin-like division or the complement-like division.

**Alpha-2-macroglobulin-like proteins:**

The alpha-2-macroglobulin-like proteins are large glycoproteins which act as non-specific irreversible proteinase inhibitors and which are found in the plasma of vertebrates, in the hemolymph of invertebrates such as lobster and in bird and reptile egg whites (Sottrup-Jensen L *et al*, 1989, J Biol Chem. 264(20):11539-42).

In humans, alpha-2-macroglobulin-like proteins include human alpha-2-macroglobulin and human Pregnancy Zone Protein (PZP). These proteinase inhibitors play a vital role in the clearance of proteinases from the circulation and in regulating proteinase activity in fibrinolysis, coagulation and complement activation.

Human alpha-2-macroglobulin is the largest known proteinase inhibitor ( $M_r=720,000$ ). It is a homotetramer formed by two protomeric units, each of which contains two 180-kDa subunits linked by two disulfide bridges. (Qazi *et al*, 1998, J Biol Chem. 273(15):8987-93. Each subunit of human alpha-2-macroglobulin has a bait region of approximately 40 amino acid residues, an internal thiol ester bond and a receptor-binding domain. Cleavage of the bait region by an attacking proteinase causes activation and cleavage of the internal thiol ester bond. This triggers major structural changes in the alpha-2-macroglobulin, known as the "mouse trap mechanism", which result in the proteinase being entrapped by and covalently linked to the alpha-2-macroglobulin. Formation of this alpha-2-macroglobulin-proteinase complex results in the exposure of the receptor binding domain of the alpha-2-macroglobulin and engagement of the receptor binding domain by cell-membrane receptors permits clearance of the alpha-2-macroglobulin-proteinase complex from circulation, via endocytosis. In contrast to the mode of inhibition of all other natural proteinase inhibitors, the entrapped proteinase retains its catalytic activity. Although inaccessible to its target proteins, the entrapped proteinase may react with small substrates and inhibitors (Qazi *et al*, 1998, *supra*).

Rats contain at least three different alpha-2-macroglobulin-like proteins, alpha 2-macroglobulin, alpha 1 inhibitor III and alpha 1-macroglobulin, which act as broad range proteinase inhibitors using a similar mechanism to known human alpha-2-macroglobulin-like proteins (Eggertsen G *et al* 1991). Chickens contain an alpha-2-macroglobulin-like protein, ovostatin, in egg white. Ovostatin differs from the alpha-2-macroglobulin-like proteins found in humans and rats in that it is more substrate specific, inhibiting only metalloproteinases stoichiometrically. Furthermore, ovostatin lacks the thiol ester bond

that other family members possess so that its mechanism of action does not involve establishing a covalent linkage between ovastatin and the proteinase (Nagase *et al*, 1986, J Biol Chem. 261(3):1421-6.)

Alpha-2-macroglobulin-like proteinase inhibitors have been implicated in a number of diseases in humans. Alterations in the serum level of human alpha-2-macroglobulin and pregnancy zone protein have been suggested to be indicative of a number of diseases and disorders. Decreased alpha-2-macroglobulin concentration typically results from enhanced clearance of alpha-2-macroglobulin-proteinase complex and occurs in states of increased proteolytic activity, such as pancreatitis. Increased serum alpha-2-macroglobulin is frequently seen in nephrotic conditions (Petersen, 1993, Dan Med Bull. 40(4):409-46), and it has been shown that proteinase inhibitory activity is lower in patients with idiopathic nephrotic syndrome (Asami *et al*, 1996, Nephron 72(4):512-7). Increased serum levels of Pregnancy Zone Protein may be an indication of threatened abortion, as well as trophoblastic diseases and gynaecological tumours (Teng H *et al*, 1994, Chin Med J (Engl) 107(12):910-4). Furthermore, pregnancy zone protein and alpha-2-macroglobulin are both able to interact with *Trypanosoma cruzi* proteinases and it has been suggested that they could prevent or minimize harmful action of *T. cruzi* proteinases, such as cruzipain, on human host molecules and regulate parasite functions controlled by cruzipain (Ramos *et al*, 2002, Exp Parasitol. 100(2):121-30.)

A number of studies have linked a valine to isoleucine (Val1000Ile) polymorphism in human alpha-2-macroglobulin with argyrophilic grain disease (AGD), a neurodegenerative disorder of the aged human brain associated with the formation of abnormal tau protein in specific neurones and macroglial cells (Ghebremedhin E *et al*, 2002, Neuropathol Appl Neurobiol (4):308-13), Alzheimer's Disease and Parkinson's Disease (Tang G *et al*, 2002, Neurosci Lett 328(2):195-7; Zappia *et al*, 2002, Neurology 59(5):756-8). However, other studies have suggested that this polymorphism does not represent a risk factor for Parkinson's Disease (Nicoletti G *et al*, 2002, Neurosci Lett 328(1):65-7).

### **Complement-like proteins**

Complement components C3, C4 and C5 are focal points in the complement system, each interacting with numerous other components during complement activation, regulation, and receptor-mediated functions. These proteins are involved in a wide variety of

biological activities such as in innate response and host defence (Fritzinger *et al*, 1992, J. Immunol. 149: 3554-3562).

C3, C4 and C5 belong to the alpha macroglobulin family but contain specific features that are not present in alpha-2-macroglobulin-like proteins, including an anaphylatoxin domain, a C-terminal netrin (NTR) domain and stretches of basic residues for proteolytic processing to form multiple chain structures. (Martinez *et al*, 2001, Front Biosci 1; 6:D904-13). Activation of C3, C4 and C5 leads to enzymatic cleavage producing fragments C3a, C4a and C5a (Ogata *et al*, 1989, J. Biol. Chem. 264: 16565-16572). Each a-fragment forms a distinct structural domain of approximately 76 amino acids, coded for by a single exon within its respective complement protein gene. (Ogata *et al*, 1989, *supra*; Gennaro *et al*, 1986, Eur. J. Biochem. 155: 77-86). The fragments are highly hydrophilic, with a mainly helical structure held together by 3 disulphide bridges (Gennaro *et al*, 1986, *supra*). The fragments are anaphylatoxins, causing smooth muscle contraction, histamine release from mast cells, and enhanced vascular permeability (Gennaro *et al*, 1986, *supra*). They also mediate chemotaxis, inflammation, and generation of cytotoxic oxygen radicals (Kohl, 2001, Mol Immunol 38(2-3):175-87).

The C-terminal netrin (NTR) domain of C3, C4 and C5 (also known as the C345C module) is also found in other proteins such as the netrins and tissue inhibitor metalloproteases (TIMPs). The functional role of NTR domains is generally unknown with the exception of TIMPs, where the NTR domain is known to be a binding site for the metalloproteinase and C5, where the NTR domain is known to be a binding site for the CP convertase, an enzyme responsible for proteolytic processing (Sandoval *et al*, 2000, J Immunol 165(2):1066-73).

Complement proteins and C3, C4 and C5 in particular, have been implicated in a variety of diseases and disorders. Generally, the anaphylatoxins formed by cleavage of C3, C4 and C5 may play a role in sepsis, immune complex disease, delayed type hypersensitivity and asthma. More specifically, C5a has been found to exert an anti-inflammatory effect in acute pancreatitis and associated lung injury (Bhatia M *et al*, 2001, Am J Physiol Gastrointest Liver Physiol 280(5):G974-8) but to induce a chronic microglia-mediated focal inflammatory response in Alzheimer's Disease (O'Barr S *et al*, 2000, J Neuroimmunol 109(2):87). Complement proteins also appear to play a role in the pathophysiology of ischaemic heart diseases and it has been suggested that complement inhibitors might be used in the treatment of this disease (Shernan SK *et al*, 2001, BioDrugs



15(9):595-607). It has also been suggested that the C4 genes may be the disease-predisposing genes connected to susceptibility to Psoriasis vulgaris (Cislo *et al*, 2002, Immunol Lett 2002 80(3):145-9).

### CD109

5 CD109 is a new member of the alpha macroglobulin family whose function remains largely unknown (Lin *et al.*, 2002, Blood 99(5):1683-91). In terms of sequence similarity, it appears to be closely related to alpha-2-macroglobulin-like proteins and more distantly related to C3 and C4 proteins. However, CD109 differs from typical alpha-2-macroglobulin-like proteinase inhibitors in several respects. Unlike alpha-2-macroglobulin-like proteinase inhibitors which generally exist as tetramers, CD109 exists  
10 as a monomer. CD109 does not contain a receptor binding domain present in alpha-2-macroglobulin-like proteinase inhibitors (Nielsen *et al*, 1996, J Biol Chem 271(22):12909-12) and unlike alpha-2-macroglobulin-like proteinase inhibitors, CD109 is membrane bound through a GPI linker. Furthermore, although CD109 contains an thioester bond  
15 similar to that found in alpha-2-macroglobulin-like proteinase inhibitors, its chemical reactivity resembles that of complement proteins. It is therefore unclear which division this novel member of the alpha macroglobulin family belongs to.

Increasing knowledge of the alpha macroglobulin family is of extreme importance in increasing the understanding of the underlying pathways that lead to the disease states and  
20 associated disease states mentioned above, and in developing more effect gene and/or drug therapies to treat these disorders. In particular, increasing knowledge of the alpha-2-macroglobulin-like proteinase inhibitors is of importance in understanding the disease states in which these proteins are implicated and developing therapies to treat these disorders.

25

### THE INVENTION

The invention is based on the discovery that the INSP087 and INSP088 proteins function as alpha macroglobulins and in particular as alpha-2-macroglobulin-like proteinase inhibitors.

30 In one embodiment of the first aspect of the invention, there is provided a polypeptide which:

- (i) comprises the amino acid sequence as recited in SEQ ID NO:2, SEQ ID NO:4,

SEQ ID NO: 6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14,  
 SEQ ID NO:16, SEQ IS NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24,  
 SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO: 32, SEQ ID NO:34,  
 SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO: 42, SEQ ID NO:44,  
 5 SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO:54,  
 SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64,  
 SEQ ID NO:66 and/or SEQ ID NO:68;

(ii) is a fragment thereof which functions as an alpha-2-macroglobulin-like proteinase inhibitor, or has an antigenic determinant in common with a polypeptide according  
 10 to (i); or

(iii) is a functional equivalent of (i) or (ii).

Preferably, the polypeptide according to this first embodiment of this first aspect of the invention:

- (i) comprises the amino acid sequence as recited in SEQ ID NO:68;
- 15 (ii) is a fragment thereof which functions as an alpha-2-macroglobulin-like proteinase inhibitor, or has an antigenic determinant in common with a polypeptide according to (i); or
- (iii) is a functional equivalent of (i) or (ii).

According to a second embodiment of this first aspect of the invention, there is provided a  
 20 polypeptide which:

- (i) consists of the amino acid sequence as recited in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ IS NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO: 32, SEQ ID NO:34,  
 25 SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO: 42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66 and/or SEQ ID NO:68;
- (ii) is a fragment thereof which functions as an alpha-2-macroglobulin-like proteinase inhibitor, or has an antigenic determinant in common with a polypeptide according  
 30 to (i); or

(iii) is a functional equivalent of (i) or (ii).

The polypeptide having the sequence recited in SEQ ID NO:2 is referred to hereafter as "the INSP087 exon 1 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:4 is referred to hereafter as "the INSP087 exon 2 polypeptide". The polypeptide  
5 having the sequence recited in SEQ ID NO:6 is referred to hereafter as "the INSP087 exon 3 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:8 is referred to hereafter as "the INSP087 exon 4 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:10 is referred to hereafter as "the INSP087 exon 5 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:12 is referred to hereafter as "the  
10 INSP087 exon 6 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:14 is referred to hereafter as "the INSP087 exon 7 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:16 is referred to hereafter as "the INSP087 exon 8 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:18 is referred to hereafter as "the INSP087 exon 9 polypeptide". The polypeptide having the  
15 sequence recited in SEQ ID NO:20 is referred to hereafter as "the INSP087 exon 10 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:22 is referred to hereafter as "the INSP087 exon 11 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:24 is referred to hereafter as "the INSP087 exon 12 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:26 is referred to hereafter as  
20 "the INSP087 exon 13 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:28 is referred to hereafter as "the INSP087 exon 14 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:30 is referred to hereafter as "the INSP087 exon 15 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:32 is referred to hereafter as "the INSP087 exon 16 polypeptide". The polypeptide having the  
25 sequence recited in SEQ ID NO:34 is referred to hereafter as "the INSP087 exon 17 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:36 is referred to hereafter as "the INSP087 exon 18 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:38 is referred to hereafter as "the INSP087 exon 19 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:40 is referred to hereafter as  
30 "the INSP087 exon 20 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:42 is referred to hereafter as "the INSP087 exon 21 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:44 is referred to hereafter as "the INSP087 exon 22 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:46 is

referred to hereafter as "the INSP087 exon 23 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:48 is referred to hereafter as "the INSP087 exon 24 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:50 is referred to hereafter as "the INSP087 exon 25 polypeptide". The polypeptide having the sequence  
5 recited in SEQ ID NO:52 is referred to hereafter as "the INSP087 exon 26 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:54 is referred to hereafter as "the INSP087 exon 27 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:56 is referred to hereafter as "the INSP087 exon 28 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:58 is referred to hereafter as "the INSP087  
10 exon 29 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:60 is referred to hereafter as "the INSP087 30 exon polypeptide". The polypeptide having the sequence recited in SEQ ID NO:62 is referred to hereafter as "the INSP087 exon 31 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:64 is referred to hereafter as "the INSP087 exon 32 polypeptide". The polypeptide having the sequence  
15 recited in SEQ ID NO:66 is referred to hereafter as "the INSP087 exon 33 polypeptide".

The polypeptide having the sequence recited in SEQ ID NO: 68 is referred to hereafter as "the INSP087 partial polypeptide". It is anticipated that the INSP087 partial polypeptide forms part of a larger polypeptide. As there is no methionine start codon at the start of the INSP087 partial polypeptide (SEQ ID NO:68), it is considered likely that the full-length  
20 INSP087 polypeptide contains additional amino acids 5' of the partial sequence recited in SEQ ID NO:68.

The term "INSP087 polypeptides" as used herein includes polypeptides comprising or consisting of the INSP087 exon 1 polypeptide, the INSP087 exon 2 polypeptide, the INSP087 exon 3 polypeptide, the INSP087 exon 4 polypeptide, the INSP087 exon 5  
25 polypeptide, the INSP087 exon 6 polypeptide, the INSP087 exon 7 polypeptide, the INSP087 exon 8 polypeptide, the INSP087 exon 9 polypeptide, the INSP087 exon 10 polypeptide, the INSP087 exon 11 polypeptide, the INSP087 exon 12 polypeptide, the INSP087 exon 13 polypeptide, the INSP087 exon 14 polypeptide, the INSP087 exon 15 polypeptide, the INSP087 exon 16 polypeptide, the INSP087 exon 17 polypeptide, the  
30 INSP087 exon 18 polypeptide, the INSP087 exon 19 polypeptide, the INSP087 exon 20 polypeptide, the INSP087 exon 21 polypeptide, the INSP087 exon 22 polypeptide, the INSP087 exon 23 polypeptide, the INSP087 exon 24 polypeptide, the INSP087 exon 25 polypeptide, the INSP087 exon 26 polypeptide, the INSP087 exon 27 polypeptide, the

INSP087 exon 28 polypeptide, the INSP087 exon 29 polypeptide, the INSP087 exon 30 polypeptide, the INSP087 exon 31 polypeptide, the INSP087 exon 32 polypeptide, the INSP087 exon 33 polypeptide and the INSP087 partial polypeptide.

In a third embodiment of the first aspect of the invention, there is provided a polypeptide  
5 which:

- (i) comprises the amino acid sequence as recited in SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO: 74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ IS NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO: 100, SEQ ID  
10 NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO: 110, and/or SEQ ID NO:112;
- (ii) is a fragment thereof which functions as an alpha-2-macroglobulin-like proteinase inhibitor, or has an antigenic determinant in common with a polypeptide according to (i); or
- 15 (iii) is a functional equivalent of (i) or (ii).

Preferably, the polypeptide according to this third embodiment of the first aspect of the invention:

- (i) comprises the amino acid sequence as recited in SEQ ID NO:112;
- (ii) is a fragment thereof which functions as an alpha-2-macroglobulin-like proteinase  
20 inhibitor, or has an antigenic determinant in common with a polypeptide according to (i); or
- (iii) is a functional equivalent of (i) or (ii).

In a fourth embodiment of the first aspect of the invention, there is provided a polypeptide which:

- 25 (i) consists of the amino acid sequence as recited in SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO: 74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ IS NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO: 100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO: 110,  
30 and/or SEQ ID NO:112;

- (ii) is a fragment thereof which functions as an alpha-2-macroglobulin-like proteinase inhibitor, or has an antigenic determinant in common with a polypeptide according to (i); or
  - (iii) is a functional equivalent of (i) or (ii).
- 5 The polypeptide having the sequence recited in SEQ ID NO:70 is referred to hereafter as "the INSP088 exon 1 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:72 is referred to hereafter as "the INSP088 exon 2 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:74 is referred to hereafter as "the INSP088 exon 3 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:76 is referred to hereafter as "the INSP088 exon 4 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:78 is referred to hereafter as "the INSP088 exon 5 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:80 is referred to hereafter as "the INSP088 exon 6 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:82 is referred to hereafter as "the INSP088 exon 7 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:84 is referred to hereafter as "the INSP088 exon 8 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:86 is referred to hereafter as "the INSP088 exon 9 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:88 is referred to hereafter as "the INSP088 exon 10 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:90 is referred to hereafter as "the INSP088 exon 11 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:92 is referred to hereafter as "the INSP088 exon 12 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:94 is referred to hereafter as "the INSP088 exon 13 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:96 is referred to hereafter as "the INSP088 exon 14 polypeptide".
- 25 The polypeptide having the sequence recited in SEQ ID NO:98 is referred to hereafter as "the INSP088 exon 15 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:100 is referred to hereafter as "the INSP088 exon 16 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:102 is referred to hereafter as "the INSP088 exon 17 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:104 is referred to hereafter as "the INSP088 exon 18 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:106 is referred to hereafter as "the INSP088 exon 19 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:108 is referred to hereafter as "the INSP088 exon 20 polypeptide". The polypeptide having the sequence

recited in SEQ ID NO:110 is referred to hereafter as "the INSP088 exon 21 polypeptide".

The polypeptide having the sequence recited in SEQ ID NO: 112 is referred to hereafter as "the INSP088 partial polypeptide". It is anticipated that the INSP088 partial polypeptide forms part of a larger polypeptide. As there is no methionine start codon at the start of the  
5 INSP087 partial polypeptide (SEQ ID NO:112), it is considered likely that the full-length INSP087 polypeptide contains additional amino acids 5' of the partial sequence recited in SEQ ID NO:112.

The term "INSP088 polypeptides" as used herein includes polypeptides comprising or consisting of the INSP088 exon 1 polypeptide, the INSP088 exon 2 polypeptide, the  
10 INSP088 exon 3 polypeptide, the INSP088 exon 4 polypeptide, the INSP088 exon 5 polypeptide, the INSP088 exon 6 polypeptide, the INSP088 exon 7 polypeptide, the INSP088 exon 8 polypeptide, the INSP088 exon 9 polypeptide, the INSP088 exon 10 polypeptide, the INSP088 exon 11 polypeptide, the INSP088 exon 12 polypeptide, the INSP088 exon 13 polypeptide, the INSP088 exon 14 polypeptide, the INSP088 exon 15  
15 polypeptide, the INSP088 exon 16 polypeptide, the INSP088 exon 17 polypeptide, the INSP088 exon 18 polypeptide, the INSP088 exon 19 polypeptide, the INSP088 exon 20 polypeptide, the INSP088 exon 21 polypeptide and the INSP088 partial polypeptide.

By "functions as an alpha-2-macroglobulin-like proteinase inhibitor" we refer to polypeptides that comprise amino acid sequence or structural features that can be identified  
20 as conserved features within alpha-2-macroglobulin-like proteinase inhibitors, such that the polypeptide's interaction with a proteinase is not substantially affected detrimentally in comparison to the function of the full length wild type polypeptide. Examples of assays which may be used to determine the biological activity of a polypeptide of the invention are described in Example 6.

25 The receptor binding domain of INSP087 has been identified as being located within amino acid residues 1263-1357 of the INSP087 partial polypeptide (SEQ ID NO:68). The amino acid sequence of the receptor binding domain of INSP087 is recited herein as SEQ ID NO:113. The receptor binding domain of INSP087 has been cloned, as described in Example 3. The sequence of the cloned receptor binding domain of INSP087, which  
30 differs from by just one amino acid residue from SEQ ID NO:113, is given herein as SEQ ID NO:115. A preferred polypeptide fragment according to the first aspect of the invention therefore comprises or consists of SEQ ID NO:113 or SEQ ID NO:115 or is a functional

equivalent thereof. It is considered highly likely, however that the receptor binding domain will fold correctly and show biological activity if additional residues C terminal and/or N terminal of these boundaries in the polypeptide sequence are included in the polypeptide fragment. For example, an additional 5, 10, 20, 30, 40, 50, 100 or even as many as 200 amino acid residues from the INSP087 polypeptide sequence, or from a homologous sequence, may be included at either or both the C terminal and/or N terminal of the boundaries of the receptor binding domain, without prejudicing the ability of the polypeptide fragment to fold correctly and exhibit receptor binding domain activity. Extensions as large as 100 or 200 residues may be necessary due to the presence of large loops between secondary structural elements.

For truncated variants of the INSP087 receptor binding domain, one or a few amino acid residues (for example, 2, 3, 4, 5, 10, 15, 20, 25, 30 or more) may be deleted at either or both the C terminus or the N terminus of the domain without prejudicing biological activity.

The receptor binding domain of INSP088 has been identified as being located within amino acid residues 803-894 of the INSP088 partial polypeptide (SEQ ID NO:112). A preferred polypeptide fragment according to the first aspect of the invention therefore comprises or consists of amino acid residues 803-894 of SEQ ID NO:112 or is a functional equivalent thereof. The amino acid sequence of the receptor binding domain of INSP088 is recited herein SEQ ID NO:117. The receptor binding domain of INSP087 has been cloned, as described in Example 4, confirming the sequence is correct. A preferred polypeptide fragment according to the first aspect of the invention therefore comprises or consists of SEQ ID NO:117 or is a functional equivalent thereof. It is considered highly likely, however that the receptor binding domain will fold correctly and show biological activity if additional residues C terminal and/or N terminal of these boundaries in the polypeptide sequence are included in the polypeptide fragment. For example, an additional 5, 10, 20, 30, 40, 50, 100 or even as many as 200 amino acid residues from the INSP088 polypeptide sequence, or from a homologous sequence, may be included at either or both the C terminal and/or N terminal of the boundaries of the receptor binding domain, without prejudicing the ability of the polypeptide fragment to fold correctly and exhibit receptor binding domain activity. Extensions as large as 100 or 200 residues may be necessary due to the presence of large loops between secondary structural elements.



For truncated variants of the INSP088 receptor binding domain, one or a few amino acid residues (for example, 2, 3, 4, 5, 10, 15, 20, 25, 30 or more) may be deleted at either or both the C terminus or the N terminus of the domain without prejudicing biological activity.

- 5 The macroglobulin domain of INSP088 has been identified as being located within amino acid residues 163-894 of the INSP088 partial polypeptide (SEQ ID NO:112). This amino acid sequence is recited herein as SEQ ID NO:119. A preferred polypeptide fragment according to the first aspect of the invention therefore comprises or consists of amino acid of SEQ ID NO:119 or is a functional equivalent thereof. It is considered highly likely,  
10 however that the macroglobulin domain will fold correctly and show biological activity if additional residues C terminal and/or N terminal of these boundaries in the polypeptide sequence are included in the polypeptide fragment. For example, an additional 5, 10, 20, 30, 40, 50, 100 or even as many as 200 amino acid residues from the INSP088 polypeptide sequence, or from a homologous sequence, may be included at either or both the C  
15 terminal and/or N terminal of the boundaries of the macroglobulin domain, without prejudicing the ability of the polypeptide fragment to fold correctly and exhibit macroglobulin domain activity. Extensions as large as 100 or 200 residues may be necessary due to the presence of large loops between secondary structural elements.

For truncated variants of the INSP088 macroglobulin domain, one or a few amino acid  
20 residues (for example, 2, 3, 4, 5, 10, 15, 20, 25, 30 or more) may be deleted at either or both the C terminus or the N terminus of the domain without prejudicing biological activity.

A further preferred polypeptide fragment according to the first aspect of the invention comprises or consists of SEQ ID NO:121, a fragment of the INSP088 partial polypeptide  
25 including the receptor binding domain and the macroglobulin domain, cloned from human tissue as described in Example 5.

In a second aspect, the invention provides a purified nucleic acid molecule which encodes a polypeptide of the first aspect of the invention.

In a first embodiment of this aspect of the invention, the purified nucleic acid molecule  
30 comprises the nucleic acid sequence as recited in SEQ ID NO:1 (encoding the INSP087 exon 1 polypeptide), SEQ ID NO:3 (encoding the INSP087 exon 2 polypeptide), SEQ ID

NO:5 (encoding the INSP087 exon 3 polypeptide), SEQ ID NO:7 (encoding the INSP087 exon 4 polypeptide), SEQ ID NO:9 (encoding the INSP087 exon 5 polypeptide), SEQ ID NO:11 (encoding the INSP087 exon 6 polypeptide), SEQ ID NO:13 (encoding the INSP087 exon 7 polypeptide), SEQ ID NO:15 (encoding the INSP087 exon 8 polypeptide), SEQ ID NO:17 (encoding the INSP087 exon 9 polypeptide), SEQ ID NO:19 (encoding the INSP087 exon 10 polypeptide), SEQ ID NO:21 (encoding the INSP087 exon 11 polypeptide), SEQ ID NO:23 (encoding the INSP087 exon 12 polypeptide), SEQ ID NO:25 (encoding the INSP087 exon 13 polypeptide), SEQ ID NO:27 (encoding the INSP087 exon 14 polypeptide), SEQ ID NO:29 (encoding the INSP087 exon 15 polypeptide), SEQ ID NO:31 (encoding the INSP087 exon 16 polypeptide), SEQ ID NO:33 (encoding the INSP087 exon 17 polypeptide), SEQ ID NO:35 (encoding the INSP087 exon 18 polypeptide), SEQ ID NO:37 (encoding the INSP087 exon 19 polypeptide), SEQ ID NO:39 (encoding the INSP087 exon 20 polypeptide), SEQ ID NO:41 (encoding the INSP087 exon 21 polypeptide), SEQ ID NO:43 (encoding the INSP087 exon 22 polypeptide), SEQ ID NO:45 (encoding the INSP087 exon 23 polypeptide), SEQ ID NO:47 (encoding the INSP087 exon 24 polypeptide), SEQ ID NO:49 (encoding the INSP087 exon 25 polypeptide), SEQ ID NO:51 (encoding the INSP087 exon 26 polypeptide), SEQ ID NO:53 (encoding the INSP087 exon 27 polypeptide), SEQ ID NO:55 (encoding the INSP087 exon 28 polypeptide), SEQ ID NO:57 (encoding the INSP087 exon 29 polypeptide), SEQ ID NO:59 (encoding the INSP087 exon 30 polypeptide), SEQ ID NO:61 (encoding the INSP087 exon 31 polypeptide), SEQ ID NO:63 (encoding the INSP087 exon 32 polypeptide), SEQ ID NO:65 (encoding the INSP087 exon 33 polypeptide), SEQ ID NO:67 (encoding the INSP087 partial polypeptide) or is a redundant equivalent or fragment of any one of these sequences.

The invention further provides that the purified nucleic acid molecule consists of the nucleic acid sequence as recited in SEQ ID NO:1 (encoding the INSP087 exon 1 polypeptide), SEQ ID NO:3 (encoding the INSP087 exon 2 polypeptide), SEQ ID NO:5 (encoding the INSP087 exon 3 polypeptide), SEQ ID NO:7 (encoding the INSP087 exon 4 polypeptide), SEQ ID NO:9 (encoding the INSP087 exon 5 polypeptide), SEQ ID NO:11 (encoding the INSP087 exon 6 polypeptide), SEQ ID NO:13 (encoding the INSP087 exon 7 polypeptide), SEQ ID NO:15 (encoding the INSP087 exon 8 polypeptide), SEQ ID NO:17 (encoding the INSP087 exon 9 polypeptide), SEQ ID NO:19 (encoding the

INSP087 exon 10 polypeptide), SEQ ID NO:21 (encoding the INSP087 exon 11 polypeptide), SEQ ID NO:23 (encoding the INSP087 exon 12 polypeptide), SEQ ID NO:25 (encoding the INSP087 exon 13 polypeptide), SEQ ID NO:27 (encoding the INSP087 exon 14 polypeptide), SEQ ID NO:29 (encoding the INSP087 exon 15 polypeptide), SEQ ID NO:31 (encoding the INSP087 exon 16 polypeptide), SEQ ID NO:33 (encoding the INSP087 exon 17 polypeptide), SEQ ID NO:35 (encoding the INSP087 exon 18 polypeptide), SEQ ID NO:37 (encoding the INSP087 exon 19 polypeptide), SEQ ID NO:39 (encoding the INSP087 exon 20 polypeptide), SEQ ID NO:41 (encoding the INSP087 exon 21 polypeptide), SEQ ID NO:43 (encoding the INSP087 exon 22 polypeptide), SEQ ID NO:45 (encoding the INSP087 exon 23 polypeptide), SEQ ID NO:47 (encoding the INSP087 exon 24 polypeptide), SEQ ID NO:49 (encoding the INSP087 exon 25 polypeptide), SEQ ID NO:51 (encoding the INSP087 exon 26 polypeptide), SEQ ID NO:53 (encoding the INSP087 exon 27 polypeptide), SEQ ID NO:55 (encoding the INSP087 exon 28 polypeptide), SEQ ID NO:57 (encoding the INSP087 exon 29 polypeptide), SEQ ID NO:59 (encoding the INSP087 exon 30 polypeptide), SEQ ID NO:61 (encoding the INSP087 exon 31 polypeptide), SEQ ID NO:63 (encoding the INSP087 exon 32 polypeptide), SEQ ID NO:65 (encoding the INSP087 exon 33 polypeptide), SEQ ID NO:67 (encoding the INSP087 partial polypeptide) or is a redundant equivalent or fragment of any one of these sequences.

In a second embodiment of this aspect of the invention, the purified nucleic acid molecule comprises the nucleic acid sequence as recited in SEQ ID NO:69 (encoding the INSP088 exon 1 polypeptide), SEQ ID NO:71 (encoding the INSP088 exon 2 polypeptide), SEQ ID NO:73 (encoding the INSP088 exon 3 polypeptide), SEQ ID NO:75 (encoding the INSP088 exon 4 polypeptide), SEQ ID NO:77 (encoding the INSP088 exon 5 polypeptide), SEQ ID NO:79 (encoding the INSP088 exon 6 polypeptide), SEQ ID NO:81 (encoding the INSP088 exon 7 polypeptide), SEQ ID NO:83 (encoding the INSP088 exon 8 polypeptide), SEQ ID NO:85 (encoding the INSP088 exon 9 polypeptide), SEQ ID NO:87 (encoding the INSP088 exon 10 polypeptide), SEQ ID NO:89 (encoding the INSP088 exon 11 polypeptide), SEQ ID NO:91 (encoding the INSP088 exon 12 polypeptide), SEQ ID NO:93 (encoding the INSP088 exon 13 polypeptide), SEQ ID NO:95 (encoding the INSP088 exon 14 polypeptide), SEQ ID NO:97 (encoding the INSP088 exon 15 polypeptide), SEQ ID NO:99 (encoding the INSP088 exon 16

polypeptide), SEQ ID NO:101 (encoding the INSP088 exon 17 polypeptide), SEQ ID NO:103 (encoding the INSP088 exon 18 polypeptide), SEQ ID NO:105 (encoding the INSP088 exon 19 polypeptide), SEQ ID NO:107 (encoding the INSP088 exon 20 polypeptide), SEQ ID NO:109 (encoding the INSP088 exon 21 polypeptide), SEQ ID NO:111 (encoding the INSP088 polypeptide) or is a redundant equivalent or fragment of any one of these sequences.

The invention further provides that the purified nucleic acid molecule consists of the nucleic acid sequence as recited in SEQ ID NO:69 (encoding the INSP088 exon 1 polypeptide), SEQ ID NO:71 (encoding the INSP088 exon 2 polypeptide), SEQ ID NO:73 (encoding the INSP088 exon 3 polypeptide), SEQ ID NO:75 (encoding the INSP088 exon 4 polypeptide), SEQ ID NO:77 (encoding the INSP088 exon 5 polypeptide), SEQ ID NO:79 (encoding the INSP088 exon 6 polypeptide), SEQ ID NO:81 (encoding the INSP088 exon 7 polypeptide), SEQ ID NO:83 (encoding the INSP088 exon 8 polypeptide), SEQ ID NO:85 (encoding the INSP088 exon 9 polypeptide), SEQ ID NO:87 (encoding the INSP088 exon 10 polypeptide), SEQ ID NO:89 (encoding the INSP088 exon 11 polypeptide), SEQ ID NO:91 (encoding the INSP088 exon 12 polypeptide), SEQ ID NO:93 (encoding the INSP088 exon 13 polypeptide), SEQ ID NO:95 (encoding the INSP088 exon 14 polypeptide), SEQ ID NO:97 (encoding the INSP088 exon 15 polypeptide), SEQ ID NO:99 (encoding the INSP088 exon 16 polypeptide), SEQ ID NO:101 (encoding the INSP088 exon 17 polypeptide), SEQ ID NO:103 (encoding the INSP088 exon 18 polypeptide), SEQ ID NO:105 (encoding the INSP088 exon 19 polypeptide), SEQ ID NO:107 (encoding the INSP088 exon 20 polypeptide), SEQ ID NO:109 (encoding the INSP088 exon 21 polypeptide), SEQ ID NO:111 (encoding the INSP088 polypeptide) or is a redundant equivalent or fragment of any one of these sequences.

Human ESTs BE144308 (sourced from head and neck tissue) and AK057908 (sourced from brain tissue) are specifically excluded from the scope of this aspect of the invention.

Preferred nucleic acid fragments according to the invention include fragments encoding the receptor binding domain of the INSP087 partial polypeptide and fragments encoding the receptor binding domain and/or macroglobulin domain of the INSP088 partial polypeptide. Preferred nucleic acid fragments comprise or consist of nucleotides SEQ ID NO:114 or SEQ ID NO:116 (encoding the INSP087 receptor binding domain), SEQ ID NO:118

(encoding the INSP088 receptor binding domain) or SEQ ID NO:120 (encoding the INSP088 macroglobulin domain). A further preferred nucleic acid fragment according to the invention comprises or consists of SEQ ID NO:122 (encoding the cloned INSP088 fragment having the amino acid sequence of SEQ ID NO:121).

- 5 In a third aspect, the invention provides a purified nucleic acid molecule which hybridizes under high stringency conditions with a nucleic acid molecule of the second aspect of the invention.

In a fourth aspect, the invention provides a vector, such as an expression vector, that contains a nucleic acid molecule of the second or third aspect of the invention.

- 10 In a fifth aspect, the invention provides a host cell transformed with a vector of the fourth aspect of the invention.

In a sixth aspect, the invention provides a ligand which binds specifically to alpha-2-macroglobulin-like proteinase inhibitors of the first aspect of the invention. Preferably, the ligand inhibits the function of a polypeptide of the first aspect of the invention.

- 15 Ligands to a polypeptide according to the invention may come in various forms including natural or modified substrates, enzymes, receptors, small organic molecules such as small natural or synthetic organic molecules of up to 2000Da, preferably 800Da or less, peptidomimetics, inorganic molecules, peptides, polypeptides, antibodies, structural or functional mimetics of the aforementioned.

- 20 In a seventh aspect, the invention provides a compound that is effective to alter the expression of a natural gene which encodes a polypeptide of the first aspect of the invention or to regulate the activity of a polypeptide of the first aspect of the invention.

Such compounds may be identified using the assays and screening methods disclosed herein.

- 25 A compound of the seventh aspect of the invention may either increase (agonise) or decrease (antagonise) the level of expression of the gene or the activity of the polypeptide.

Importantly, the identification of the function of the INSP087 and INSP088 polypeptides allows for the design of screening methods capable of identifying compounds that are effective in the treatment and/or diagnosis of disease. Ligands and compounds according to

- 30 the sixth and seventh aspects of the invention may be identified using such methods. Examples of suitable assays and screening methods are provided herein. These methods

are included as aspects of the present invention.

In an eighth aspect, the invention provides a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the fourth aspect of the invention, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention, for use in therapy or diagnosis of a disease in which alpha-2-macroglobulin-like proteinase inhibitors are implicated. Such diseases and disorders may include reproductive disorders, cell proliferative disorders, including neoplasm, melanoma, lung, colorectal, breast, pancreas, head and neck and other solid tumours; myeloproliferative disorders, such as leukemia, non-Hodgkin lymphoma, leukopenia, thrombocytopenia, angiogenesis disorder, Kaposi's sarcoma; autoimmune/inflammatory disorders, including allergy, inflammatory bowel disease, pancreatitis, arthritis, psoriasis, psoriasis vulgaris, respiratory tract inflammation, asthma, and organ transplant rejection; cardiovascular disorders, including hypertension, oedema, angina, atherosclerosis, thrombosis, sepsis, shock, reperfusion injury, and ischemia, particularly ischemic heart disease; neurological disorders including central nervous system disease, Alzheimer's disease, brain injury, Parkinson's disease, amyotrophic lateral sclerosis, and pain; developmental disorders; metabolic disorders including diabetes mellitus, osteoporosis, and obesity, AIDS, renal disease, particularly idiopathic nephrotic syndrome; lung injury; infections including viral infection, bacterial infection, fungal infection and parasitic infection, particularly *Trypanosoma cruzi* infection and other pathological conditions. Preferably, the disease is one in which alpha-2-macroglobulin-like proteinase inhibitors are implicated. These molecules may also be used in the manufacture of a medicament for the treatment of such disorders.

In a ninth aspect, the invention provides a method of diagnosing a disease in a patient, comprising assessing the level of expression of a natural gene encoding a polypeptide of the first aspect of the invention or the activity of a polypeptide of the first aspect of the invention in tissue from said patient and comparing said level of expression or activity to a control level, wherein a level that is different to said control level is indicative of disease. Such a method will preferably be carried out *in vitro*. Similar methods may be used for monitoring the therapeutic treatment of disease in a patient, wherein altering the level of expression or activity of a polypeptide or nucleic acid molecule over the period of time towards a control level is indicative of regression of disease.

Preferably, the disease diagnosed by a method of a ninth aspect of the invention is a disease in which alpha-2-macroglobulin-like proteinase inhibitors are implicated, as described above.

A preferred method for detecting polypeptides of the first aspect of the invention  
5 comprises the steps of: (a) contacting a ligand, such as an antibody, of the sixth aspect of the invention with a biological sample under conditions suitable for the formation of a ligand-polypeptide complex; and (b) detecting said complex.

A number of different methods according to the ninth aspect of the invention exist, as the skilled reader will be aware, such as methods of nucleic acid hybridization with short  
10 probes, point mutation analysis, polymerase chain reaction (PCR) amplification and methods using antibodies to detect aberrant protein levels. Similar methods may be used on a short or long term basis to allow therapeutic treatment of a disease to be monitored in a patient. The invention also provides kits that are useful in these methods for diagnosing disease.

15 In a tenth aspect, the invention provides for the use of the polypeptides of the first aspect of the invention as alpha-2-macroglobulin-like proteinase inhibitors. Suitable uses of the polypeptides of the invention as alpha-2-macroglobulin-like proteinase inhibitors include use as a diagnostic marker for a physiological or pathological condition selected from the list given above.

20 In an eleventh aspect, the invention provides a pharmaceutical composition comprising a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the fourth aspect of the invention, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention, in conjunction with a pharmaceutically-acceptable carrier.

25 In a twelfth aspect, the present invention provides a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the fourth aspect of the invention, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention, for use in therapy or diagnosis. These molecules may also be used in the manufacture of a medicament for the treatment of  
30 a disease.

In a thirteenth aspect, the invention provides a method of treating a disease in a patient

comprising administering to the patient a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the fourth aspect of the invention, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention.

- 5 For diseases in which the expression of a natural gene encoding a polypeptide of the first aspect of the invention, or in which the activity of a polypeptide of the first aspect of the invention, is lower in a diseased patient when compared to the level of expression or activity in a healthy patient, the polypeptide, nucleic acid molecule, ligand or compound administered to the patient should be an agonist. Conversely, for diseases in which the
- 10 expression of the natural gene or activity of the polypeptide is higher in a diseased patient when compared to the level of expression or activity in a healthy patient, the polypeptide, nucleic acid molecule, ligand or compound administered to the patient should be an antagonist. Examples of such antagonists include antisense nucleic acid molecules, ribozymes and ligands, such as antibodies.
- 15 In a fourteenth aspect, the invention provides transgenic or knockout non-human animals that have been transformed to express higher, lower or absent levels of a polypeptide of the first aspect of the invention. Such transgenic animals are very useful models for the study of disease and may also be used in screening regimes for the identification of compounds that are effective in the treatment or diagnosis of such a disease. Preferably, the disease a
- 20 disease in which alpha-2-macroglobulin-like proteinase inhibitors are implicated, as described above.

A summary of standard techniques and procedures which may be employed in order to utilise the invention is given below. It will be understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors and reagents described. It is

25 also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and it is not intended that this terminology should limit the scope of the present invention. The extent of the invention is limited only by the terms of the appended claims.

Standard abbreviations for nucleotides and amino acids are used in this specification.

- 30 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA technology and



immunology, which are within the skill of those working in the art.

Such techniques are explained fully in the literature. Examples of particularly suitable texts for consultation include the following: Sambrook Molecular Cloning; A Laboratory Manual, Second Edition (1989); DNA Cloning, Volumes I and II (D.N Glover ed. 1985);  
5 Oligonucleotide Synthesis (M.J. Gait ed. 1984); Nucleic Acid Hybridization (B.D. Hames & S.J. Higgins eds. 1984); Transcription and Translation (B.D. Hames & S.J. Higgins eds. 1984); Animal Cell Culture (R.I. Freshney ed. 1986); Immobilized Cells and Enzymes (IRL Press, 1986); B. Perbal, A Practical Guide to Molecular Cloning (1984); the Methods in Enzymology series (Academic Press, Inc.), especially volumes 154 & 155; Gene  
10 Transfer Vectors for Mammalian Cells (J.H. Miller and M.P. Calos eds. 1987, Cold Spring Harbor Laboratory); Immunochemical Methods in Cell and Molecular Biology (Mayer and Walker, eds. 1987, Academic Press, London); Scopes, (1987) Protein Purification: Principles and Practice, Second Edition (Springer Verlag, N.Y.); and Handbook of Experimental Immunology, Volumes I-IV (D.M. Weir and C. C. Blackwell eds. 1986).

- 15 As used herein, the term "polypeptide" includes any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e. peptide isosteres. This term refers both to short chains (peptides and oligopeptides) and to longer chains (proteins).

The polypeptide of the present invention may be in the form of a mature protein or may be  
20 a pre-, pro- or prepro- protein that can be activated by cleavage of the pre-, pro- or prepro- portion to produce an active mature polypeptide. In such polypeptides, the pre-, pro- or prepro- sequence may be a leader or secretory sequence or may be a sequence that is employed for purification of the mature polypeptide sequence.

The polypeptide of the first aspect of the invention may form part of a fusion protein. For  
25 example, it is often advantageous to include one or more additional amino acid sequences which may contain secretory or leader sequences, pro-sequences, sequences which aid in purification, or sequences that confer higher protein stability, for example during recombinant production. Alternatively or additionally, the mature polypeptide may be fused with another compound, such as a compound to increase the half-life of the  
30 polypeptide (for example, polyethylene glycol).

Polypeptides may contain amino acids other than the 20 gene-encoded amino acids, modified either by natural processes, such as by post-translational processing or by

chemical modification techniques which are well known in the art. Among the known modifications which may commonly be present in polypeptides of the present invention are glycosylation, lipid attachment, sulphation, gamma-carboxylation, for instance of glutamic acid residues, hydroxylation and ADP-ribosylation. Other potential modifications  
5 include acetylation, acylation, amidation, covalent attachment of flavin, covalent attachment of a haeme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulphide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate,  
10 formylation, GPI anchor formation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. In fact, blockage of the amino  
15 or carboxyl terminus in a polypeptide, or both, by a covalent modification is common in naturally-occurring and synthetic polypeptides and such modifications may be present in polypeptides of the present invention.

The modifications that occur in a polypeptide often will be a function of how the polypeptide is made. For polypeptides that are made recombinantly, the nature and extent  
20 of the modifications in large part will be determined by the post-translational modification capacity of the particular host cell and the modification signals that are present in the amino acid sequence of the polypeptide in question. For instance, glycosylation patterns vary between different types of host cell.

The polypeptides of the present invention can be prepared in any suitable manner. Such  
25 polypeptides include isolated naturally-occurring polypeptides (for example purified from cell culture), recombinantly-produced polypeptides (including fusion proteins), synthetically-produced polypeptides or polypeptides that are produced by a combination of these methods.

The functionally-equivalent polypeptides of the first aspect of the invention may be  
30 polypeptides that are homologous to the INSP087 and INSP088 polypeptides. Two polypeptides are said to be "homologous", as the term is used herein, if the sequence of one of the polypeptides has a high enough degree of identity or similarity to the sequence of

the other polypeptide. "Identity" indicates that at any particular position in the aligned sequences, the amino acid residue is identical between the sequences. "Similarity" indicates that, at any particular position in the aligned sequences, the amino acid residue is of a similar type between the sequences. Degrees of identity and similarity can be readily  
5 calculated (Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York, 1988; Biocomputing. Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer,  
10 Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991).

Homologous polypeptides therefore include natural biological variants (for example, allelic variants or geographical variations within the species from which the polypeptides are derived) and mutants (such as mutants containing amino acid substitutions, insertions or deletions) of the INSP087 and INSP088 polypeptides. Such mutants may include  
15 polypeptides in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; among the basic  
20 residues Lys and Arg; or among the aromatic residues Phe and Tyr. Particularly preferred are variants in which several, i.e. between 5 and 10, 1 and 5, 1 and 3, 1 and 2 or just 1 amino acids are substituted, deleted or added in any combination. Especially preferred are silent substitutions, additions and deletions, which do not alter the properties and activities of the protein. Also especially preferred in this regard are conservative substitutions. Such  
25 mutants also include polypeptides in which one or more of the amino acid residues includes a substituent group.

Typically, greater than 30% identity between two polypeptides is considered to be an indication of functional equivalence. Preferably, functionally equivalent polypeptides of the first aspect of the invention have a degree of sequence identity with the INSP087 and  
30 INSP088 polypeptides, or with active fragments thereof, of greater than 45%. More preferred polypeptides have degrees of identity of greater than 50%, 60%. 70%. 80%, 90%, 95%, 98% or 99%, respectively.

The functionally-equivalent polypeptides of the first aspect of the invention may also be polypeptides which have been identified using one or more techniques of structural alignment. For example, the Inpharmatica Genome Threader<sup>TM</sup> technology that forms one aspect of the search tools used to generate the Biopendium search database may be used  
5 (see co-pending International Patent Application No. PCT/GB01/01105) to identify polypeptides of presently-unknown function which, while having low sequence identity as compared to the INSP087 and INSP088 polypeptides, are predicted to be alpha-2 macroglobulin-like proteinases, said method utilising a polypeptide of the first aspect of the invention, by virtue of sharing significant structural homology with the INSP087 and  
10 INSP088 polypeptide sequences. By "significant structural homology" is meant that the Inpharmatica Genome Threader<sup>TM</sup> predicts two proteins to share structural homology with a certainty of at 10% and above.

The polypeptides of the first aspect of the invention also include fragments of the INSP087 and INSP088 polypeptides and fragments of the functional equivalents of the INSP087 and  
15 INSP088 polypeptides, provided that those fragments retain alpha-2-macroglobulin-like proteinase inhibiting activity or have an antigenic determinant in common with the INSP087 and INSP088 polypeptides.

As used herein, the term "fragment" refers to a polypeptide having an amino acid sequence that is the same as part, but not all, of the amino acid sequence of the INSP087 and  
20 INSP088 polypeptides or one of its functional equivalents. The fragments should comprise at least n consecutive amino acids from the sequence and, depending on the particular sequence, n preferably is 7 or more (for example, 8, 10, 12, 14, 16, 18, 20 or more). Small fragments may form an antigenic determinant.

Fragments of the INSP087 and INSP088 polypeptides may consist of combinations of 2, 3,  
25 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 of neighbouring exon sequences in the INSP087 or INSP088 partial polypeptide sequences respectively.

Such fragments may be "free-standing", i.e. not part of or fused to other amino acids or polypeptides, or they may be comprised within a larger polypeptide of which they form a part or region. When comprised within a larger polypeptide, the fragment of the invention  
30 most preferably forms a single continuous region. For instance, certain preferred embodiments relate to a fragment having a pre - and/or pro- polypeptide region fused to the amino terminus of the fragment and/or an additional region fused to the carboxyl

terminus of the fragment. However, several fragments may be comprised within a single larger polypeptide.

The polypeptides of the present invention or their immunogenic fragments (comprising at least one antigenic determinant) can be used to generate ligands, such as polyclonal or  
5 monoclonal antibodies, that are immunospecific for the polypeptides. Such antibodies may be employed to isolate or to identify clones expressing the polypeptides of the invention or to purify the polypeptides by affinity chromatography. The antibodies may also be employed as diagnostic or therapeutic aids, amongst other applications, as will be apparent to the skilled reader.

10 The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art. As used herein, the term "antibody" refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub> and Fv, which are capable of binding to the antigenic determinant in question. Such antibodies thus bind to the polypeptides of the first  
15 aspect of the invention.

By "substantially greater affinity we mean that there is a measurable increase in the affinity for a polypeptide of the invention as compared with the affinity for known alpha-2-macroglobulin-like proteinase inhibitors.

Preferably, the affinity is at least 1.5-fold, 2-fold, 5-fold, 10-fold, 100-fold, 10<sup>3</sup>-fold, 10<sup>4</sup>-  
20 fold, 10<sup>5</sup>-fold or 10<sup>6</sup>-fold or greater for a polypeptide of the invention than for known alpha-2-macroglobulin-like proteinase inhibitors.

If polyclonal antibodies are desired, a selected mammal, such as a mouse, rabbit, goat or horse, may be immunised with a polypeptide of the first aspect of the invention. The polypeptide used to immunise the animal can be derived by recombinant DNA technology  
25 or can be synthesized chemically. If desired, the polypeptide can be conjugated to a carrier protein. Commonly used carriers to which the polypeptides may be chemically coupled include bovine serum albumin, thyroglobulin and keyhole limpet haemocyanin. The coupled polypeptide is then used to immunise the animal. Serum from the immunised animal is collected and treated according to known procedures, for example by  
30 immunoaffinity chromatography.

Monoclonal antibodies to the polypeptides of the first aspect of the invention can also be

readily produced by one skilled in the art. The general methodology for making monoclonal antibodies using hybridoma technology is well known (see, for example, Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, 77-96 in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985).

Panels of monoclonal antibodies produced against the polypeptides of the first aspect of the invention can be screened for various properties, i.e., for isotype, epitope, affinity, etc. Monoclonal antibodies are particularly useful in purification of the individual polypeptides against which they are directed. Alternatively, genes encoding the monoclonal antibodies of interest may be isolated from hybridomas, for instance by PCR techniques known in the art, and cloned and expressed in appropriate vectors.

Chimeric antibodies, in which non-human variable regions are joined or fused to human constant regions (see, for example, Liu *et al.*, *Proc. Natl. Acad. Sci. USA*, 84, 3439 (1987)), may also be of use.

The antibody may be modified to make it less immunogenic in an individual, for example by humanisation (see Jones *et al.*, *Nature*, 321, 522 (1986); Verhoeyen *et al.*, *Science*, 239, 1534 (1988); Kabat *et al.*, *J. Immunol.*, 147, 1709 (1991); Queen *et al.*, *Proc. Natl Acad. Sci. USA*, 86, 10029 (1989); Gorman *et al.*, *Proc. Natl Acad. Sci. USA*, 88, 34181 (1991); and Hodgson *et al.*, *Bio/Technology*, 9, 421 (1991)). The term "humanised antibody", as used herein, refers to antibody molecules in which the CDR amino acids and selected other amino acids in the variable domains of the heavy and/or light chains of a non-human donor antibody have been substituted in place of the equivalent amino acids in a human antibody. The humanised antibody thus closely resembles a human antibody but has the binding ability of the donor antibody.

In a further alternative, the antibody may be a "bispecific" antibody, that is an antibody having two different antigen binding domains, each domain being directed against a different epitope.

Phage display technology may be utilised to select genes which encode antibodies with binding activities towards the polypeptides of the invention either from repertoires of PCR amplified V-genes of lymphocytes from humans screened for possessing the relevant antibodies, or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can

also be improved by chain shuffling (Clackson, T. *et al.*, (1991) Nature 352, 624-628).

Antibodies generated by the above techniques, whether polyclonal or monoclonal, have additional utility in that they may be employed as reagents in immunoassays, radioimmunoassays (RIA) or enzyme-linked immunosorbent assays (ELISA). In these  
5 applications, the antibodies can be labelled with an analytically-detectable reagent such as a radioisotope, a fluorescent molecule or an enzyme.

Preferred nucleic acid molecules of the second and third aspects of the invention are those which encode a polypeptide sequence as recited in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16,  
10 SEQ IS NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO: 32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO: 42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66 SEQ ID NO:68; SEQ ID NO:70,  
15 SEQ ID NO:72, SEQ ID NO: 74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ IS NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO: 100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO: 110, and SEQ ID NO:112, and functionally equivalent polypeptides. These nucleic acid molecules may  
20 be used in the methods and applications described herein. The nucleic acid molecules of the invention preferably comprise at least n consecutive nucleotides from the sequences disclosed herein where, depending on the particular sequence, n is 10 or more (for example, 12, 14, 15, 18, 20, 25, 30, 35, 40 or more).

The nucleic acid molecules of the invention also include sequences that are complementary  
25 to nucleic acid molecules described above (for example, for antisense or probing purposes).

Nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance cDNA, synthetic DNA or genomic DNA. Such nucleic acid molecules may be obtained by cloning, by chemical synthetic  
30 techniques or by a combination thereof. The nucleic acid molecules can be prepared, for example, by chemical synthesis using techniques such as solid phase phosphoramidite chemical synthesis, from genomic or cDNA libraries or by separation from an organism.

RNA molecules may generally be generated by the *in vitro* or *in vivo* transcription of DNA sequences.

The nucleic acid molecules may be double-stranded or single-stranded. Single-stranded DNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

The term "nucleic acid molecule" also includes analogues of DNA and RNA, such as those containing modified backbones, and peptide nucleic acids (PNA). The term "PNA", as used herein, refers to an antisense molecule or an anti-gene agent which comprises an oligonucleotide of at least five nucleotides in length linked to a peptide backbone of amino acid residues, which preferably ends in lysine. The terminal lysine confers solubility to the composition. PNAs may be pegylated to extend their lifespan in a cell, where they preferentially bind complementary single stranded DNA and RNA and stop transcript elongation (Nielsen, P.E. *et al.* (1993) *Anticancer Drug Des.* 8:53-63).

A nucleic acid molecule which encodes a polypeptide of this invention may be identical to the coding sequence of one or more of the nucleic acid molecules disclosed herein.

These molecules also may have a different sequence which, as a result of the degeneracy of the genetic code, encode a polypeptide of in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, or SEQ ID NO:112.

Such nucleic acid molecules may include, but are not limited to, the coding sequence for the mature polypeptide by itself; the coding sequence for the mature polypeptide and additional coding sequences, such as those encoding a leader or secretory sequence, such as a pro-, pre- or prepro- polypeptide sequence; the coding sequence of the mature



polypeptide, with or without the aforementioned additional coding sequences, together with further additional, non-coding sequences, including non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription (including termination signals), ribosome binding and mRNA stability. The nucleic acid molecules  
5 may also include additional sequences which encode additional amino acids, such as those which provide additional functionalities.

The nucleic acid molecules of the second and third aspects of the invention may also encode the fragments or the functional equivalents of the polypeptides and fragments of the first aspect of the invention. Such a nucleic acid molecule may be a naturally-occurring  
10 variant such as a naturally-occurring allelic variant, or the molecule may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the nucleic acid molecule may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells or organisms.

Among variants in this regard are variants that differ from the aforementioned nucleic acid  
15 molecules by nucleotide substitutions, deletions or insertions. The substitutions, deletions or insertions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or insertions.

The nucleic acid molecules of the invention can also be engineered, using methods  
20 generally known in the art, for a variety of reasons, including modifying the cloning, processing, and/or expression of the gene product (the polypeptide). DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides are included as techniques which may be used to engineer the nucleotide sequences. Site-directed mutagenesis may be used to insert new restriction sites, alter  
25 glycosylation patterns, change codon preference, produce splice variants, introduce mutations and so forth.

Nucleic acid molecules which encode a polypeptide of the first aspect of the invention may be ligated to a heterologous sequence so that the combined nucleic acid molecule encodes a fusion protein. Such combined nucleic acid molecules are included within the second or  
30 third aspects of the invention. For example, to screen peptide libraries for inhibitors of the activity of the polypeptide, it may be useful to express, using such a combined nucleic acid molecule, a fusion protein that can be recognised by a commercially-available antibody. A

fusion protein may also be engineered to contain a cleavage site located between the sequence of the polypeptide of the invention and the sequence of a heterologous protein so that the polypeptide may be cleaved and purified away from the heterologous protein.

The nucleic acid molecules of the invention also include antisense molecules that are partially complementary to nucleic acid molecules encoding polypeptides of the present invention and that therefore hybridize to the encoding nucleic acid molecules (hybridization). Such antisense molecules, such as oligonucleotides, can be designed to recognise, specifically bind to and prevent transcription of a target nucleic acid encoding a polypeptide of the invention, as will be known by those of ordinary skill in the art (see, for example, Cohen, J.S., Trends in Pharm. Sci., 10, 435 (1989), Okano, J. Neurochem. 56, 560 (1991); O'Connor, J. Neurochem 56, 560 (1991); Lee *et al.*, Nucleic Acids Res 6, 3073 (1979); Cooney *et al.*, Science 241, 456 (1988); Dervan *et al.*, Science 251, 1360 (1991).

The term "hybridization" as used herein refers to the association of two nucleic acid molecules with one another by hydrogen bonding. Typically, one molecule will be fixed to a solid support and the other will be free in solution. Then, the two molecules may be placed in contact with one another under conditions that favour hydrogen bonding. Factors that affect this bonding include: the type and volume of solvent; reaction temperature; time of hybridization; agitation; agents to block the non-specific attachment of the liquid phase molecule to the solid support (Denhardt's reagent or BLOTTO); the concentration of the molecules; use of compounds to increase the rate of association of molecules (dextran sulphate or polyethylene glycol); and the stringency of the washing conditions following hybridization (see Sambrook *et al.* [*supra*]).

The inhibition of hybridization of a completely complementary molecule to a target molecule may be examined using a hybridization assay, as known in the art (see, for example, Sambrook *et al.* [*supra*]). A substantially homologous molecule will then compete for and inhibit the binding of a completely homologous molecule to the target molecule under various conditions of stringency, as taught in Wahl, G.M. and S.L. Berger (1987; Methods Enzymol. 152:399-407) and Kimmel, A.R. (1987; Methods Enzymol. 152:507-511).

"Stringency" refers to conditions in a hybridization reaction that favour the association of very similar molecules over association of molecules that differ. High stringency hybridisation conditions are defined as overnight incubation at 42°C in a solution

comprising 50% formamide, 5XSSC (150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulphate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1X SSC at approximately 65°C. Low stringency conditions involve the hybridisation  
5 reaction being carried out at 35°C (see Sambrook *et al.* [*supra*]). Preferably, the conditions used for hybridization are those of high stringency.

Preferred embodiments of this aspect of the invention are nucleic acid molecules that are at least 70% identical over their entire length to a nucleic acid molecule encoding the INSP087 or INSP088 and nucleic acid molecules that are substantially complementary to  
10 such nucleic acid molecules.

Preferably, a nucleic acid molecule according to this aspect of the invention comprises a region that is at least 80% identical over its entire length to such coding sequences, or is a nucleic acid molecule that is complementary thereto. In this regard, nucleic acid molecules at least 90%, preferably at least 95%, more preferably at least 98% or 99% identical over  
15 their entire length to the same are particularly preferred. Preferred embodiments in this respect are nucleic acid molecules that encode polypeptides which retain substantially the same biological function or activity as the INSP087 and INSP088 polypeptides.

The invention also provides a process for detecting a nucleic acid molecule of the invention, comprising the steps of: (a) contacting a nucleic probe according to the  
20 invention with a biological sample under hybridizing conditions to form duplexes; and (b) detecting any such duplexes that are formed.

As discussed additionally below in connection with assays that may be utilised according to the invention, a nucleic acid molecule as described above may be used as a hybridization probe for RNA, cDNA or genomic DNA, in order to isolate full-length  
25 cDNAs and genomic clones encoding the INSP087 and INSP088 polypeptides and to isolate cDNA and genomic clones of homologous or orthologous genes that have a high sequence similarity to the gene encoding this polypeptide.

In this regard, the following techniques, among others known in the art, may be utilised and are discussed below for purposes of illustration. Methods for DNA sequencing and  
30 analysis are well known and are generally available in the art and may, indeed, be used to practice many of the embodiments of the invention discussed herein. Such methods may employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase (US

Biochemical Corp, Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, IL), or combinations of polymerases and proof-reading exonucleases such as those found in the ELONGASE Amplification System marketed by Gibco/BRL (Gaithersburg, MD). Preferably, the sequencing process may be automated  
5 using machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, NV), the Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI Catalyst and 373 and 377 DNA Sequencers (Perkin Elmer).

One method for isolating a nucleic acid molecule encoding a polypeptide with an equivalent function to that of the INSP087 and INSP088 polypeptides is to probe a  
10 genomic or cDNA library with a natural or artificially-designed probe using standard procedures that are recognised in the art (see, for example, "Current Protocols in Molecular Biology", Ausubel *et al.* (eds). Greene Publishing Association and John Wiley Interscience, New York, 1989,1992). Probes comprising at least 15, preferably at least 30, and more preferably at least 50, contiguous bases that correspond to, or are complementary  
15 to, nucleic acid sequences from the appropriate encoding gene (SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID  
20 NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID  
25 NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109 or SEQ ID NO:111) are particularly useful probes. Such probes may be labelled with an analytically-detectable reagent to facilitate their identification. Useful reagents include, but are not limited to, radioisotopes, fluorescent dyes and enzymes that are capable of catalysing the formation of a detectable product. Using these probes, the ordinarily skilled artisan will be  
30 capable of isolating complementary copies of genomic DNA, cDNA or RNA polynucleotides encoding proteins of interest from human, mammalian or other animal sources and screening such sources for related sequences, for example, for additional members of the family, type and/or subtype.

In many cases, isolated cDNA sequences will be incomplete, in that the region encoding the polypeptide will be cut short, normally at the 5' end. Several methods are available to obtain full length cDNAs, or to extend short cDNAs. Such sequences may be extended utilising a partial nucleotide sequence and employing various methods known in the art to  
5 detect upstream sequences such as promoters and regulatory elements. For example, one method which may be employed is based on the method of Rapid Amplification of cDNA Ends (RACE; see, for example, Frohman *et al.*, PNAS USA 85, 8998-9002, 1988). Recent modifications of this technique, exemplified by the Marathon<sup>TM</sup> technology (Clontech Laboratories Inc.), for example, have significantly simplified the search for longer cDNAs.  
10 A slightly different technique, termed "restriction-site" PCR, uses universal primers to retrieve unknown nucleic acid sequence adjacent a known locus (Sarkar, G. (1993) PCR Methods Applic. 2:318-322). Inverse PCR may also be used to amplify or to extend sequences using divergent primers based on a known region (Triglia, T. *et al.* (1988) Nucleic Acids Res. 16:8186). Another method which may be used is capture PCR which  
15 involves PCR amplification of DNA fragments adjacent a known sequence in human and yeast artificial chromosome DNA (Lagerstrom, M. *et al.* (1991) PCR Methods Applic., 1, 111-119). Another method which may be used to retrieve unknown sequences is that of Parker, J.D. *et al.* (1991); Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PromoterFinder<sup>TM</sup> libraries to walk genomic DNA (Clontech,  
20 Palo Alto, CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable, in that they will contain more sequences that contain the 5' regions of genes. Use of a randomly  
25 primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

In one embodiment of the invention, the nucleic acid molecules of the present invention may be used for chromosome localisation. In this technique, a nucleic acid molecule is  
30 specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important step in the confirmatory correlation of those sequences with the gene-associated disease. Once a sequence has been mapped to a precise

chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationships between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localised by genetic linkage to a particular genomic region, any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleic acid molecule may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

The nucleic acid molecules of the present invention are also valuable for tissue localisation. Such techniques allow the determination of expression patterns of the polypeptide in tissues by detection of the mRNAs that encode them. These techniques include *in situ* hybridization techniques and nucleotide amplification techniques, such as PCR. Results from these studies provide an indication of the normal functions of the polypeptide in the organism. In addition, comparative studies of the normal expression pattern of mRNAs with that of mRNAs encoded by a mutant gene provide valuable insights into the role of mutant polypeptides in disease. Such inappropriate expression may be of a temporal, spatial or quantitative nature.

Gene silencing approaches may also be undertaken to down-regulate endogenous expression of a gene encoding a polypeptide of the invention. RNA interference (RNAi) (Elbashir, SM *et al.*, Nature 2001, 411, 494-498) is one method of sequence specific post-transcriptional gene silencing that may be employed. Short dsRNA oligonucleotides are synthesised *in vitro* and introduced into a cell. The sequence specific binding of these dsRNA oligonucleotides triggers the degradation of target mRNA, reducing or ablating target protein expression.

Efficacy of the gene silencing approaches assessed above may be assessed through the measurement of polypeptide expression (for example, by Western blotting), and at the RNA level using TaqMan-based methodologies.

The vectors of the present invention comprise nucleic acid molecules of the invention and

may be cloning or expression vectors. The host cells of the invention, which may be transformed, transfected or transduced with the vectors of the invention may be prokaryotic or eukaryotic.

The polypeptides of the invention may be prepared in recombinant form by expression of their encoding nucleic acid molecules in vectors contained within a host cell. Such expression methods are well known to those of skill in the art and many are described in detail by Sambrook *et al* (*supra*) and Fernandez & Hoeffler (1998, eds. "Gene expression systems. Using nature for the art of expression". Academic Press, San Diego, London, Boston, New York, Sydney, Tokyo, Toronto).

Generally, any system or vector that is suitable to maintain, propagate or express nucleic acid molecules to produce a polypeptide in the required host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those described in Sambrook *et al.*, (*supra*). Generally, the encoding gene can be placed under the control of a control element such as a promoter, ribosome binding site (for bacterial expression) and, optionally, an operator, so that the DNA sequence encoding the desired polypeptide is transcribed into RNA in the transformed host cell.

Examples of suitable expression systems include, for example, chromosomal, episomal and virus-derived systems, including, for example, vectors derived from: bacterial plasmids, bacteriophage, transposons, yeast episomes, insertion elements, yeast chromosomal elements, viruses such as baculoviruses, papova viruses such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, or combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, including cosmids and phagemids. Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained and expressed in a plasmid.

Particularly suitable expression systems include microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (for example, baculovirus); plant cell systems transformed with virus expression vectors (for example, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (for example, Ti or pBR322 plasmids); or animal cell systems. Cell-free translation systems can also be employed to produce the

polypeptides of the invention.

Introduction of nucleic acid molecules encoding a polypeptide of the present invention into host cells can be effected by methods described in many standard laboratory manuals, such as Davis *et al.*, Basic Methods in Molecular Biology (1986) and Sambrook *et al.*, (*supra*).

5 Particularly suitable methods include calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection (see Sambrook *et al.*, 1989 [*supra*]; Ausubel *et al.*, 1991 [*supra*]; Spector, Goldman & Leinwald, 1998). In eukaryotic cells, expression systems may either be transient (for  
10 example, episomal) or permanent (chromosomal integration) according to the needs of the system.

The encoding nucleic acid molecule may or may not include a sequence encoding a control sequence, such as a signal peptide or leader sequence, as desired, for example, for secretion of the translated polypeptide into the lumen of the endoplasmic reticulum, into  
15 the periplasmic space or into the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals. Leader sequences can be removed by the bacterial host in post-translational processing.

In addition to control sequences, it may be desirable to add regulatory sequences that allow for regulation of the expression of the polypeptide relative to the growth of the host cell.  
20 Examples of regulatory sequences are those which cause the expression of a gene to be increased or decreased in response to a chemical or physical stimulus, including the presence of a regulatory compound or to various temperature or metabolic conditions. Regulatory sequences are those non-translated regions of the vector, such as enhancers, promoters and 5' and 3' untranslated regions. These interact with host cellular proteins to  
25 carry out transcription and translation. Such regulatory sequences may vary in their strength and specificity. Depending on the vector system and host utilised, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript phagemid (Stratagene, LaJolla, CA) or pSportl<sup>TM</sup> plasmid (Gibco BRL) and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the  
30 genomes of plant cells (for example, heat shock, RUBISCO and storage protein genes) or



from plant viruses (for example, viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

An expression vector is constructed so that the particular nucleic acid coding sequence is located in the vector with the appropriate regulatory sequences, the positioning and orientation of the coding sequence with respect to the regulatory sequences being such that the coding sequence is transcribed under the "control" of the regulatory sequences, i.e., RNA polymerase which binds to the DNA molecule at the control sequences transcribes the coding sequence. In some cases it may be necessary to modify the sequence so that it may be attached to the control sequences with the appropriate orientation; i.e., to maintain the reading frame.

The control sequences and other regulatory sequences may be ligated to the nucleic acid coding sequence prior to insertion into a vector. Alternatively, the coding sequence can be cloned directly into an expression vector that already contains the control sequences and an appropriate restriction site.

For long-term, high-yield production of a recombinant polypeptide, stable expression is preferred. For example, cell lines which stably express the polypeptide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells that successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalised cell lines available from the American Type Culture Collection (ATCC) including, but not limited to, Chinese hamster ovary (CHO), HeLa, baby hamster kidney (BHK), monkey kidney (COS), C127, 3T3, BHK, HEK 293, Bowes melanoma and human hepatocellular carcinoma (for example Hep G2) cells and a number of other cell

lines.

In the baculovirus system, the materials for baculovirus/insect cell expression systems are commercially available in kit form from, inter alia, Invitrogen, San Diego CA (the "MaxBac" kit). These techniques are generally known to those skilled in the art and are described fully in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987). Particularly suitable host cells for use in this system include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells.

There are many plant cell culture and whole plant genetic expression systems known in the art. Examples of suitable plant cellular genetic expression systems include those described in US 5,693,506; US 5,659,122; and US 5,608,143. Additional examples of genetic expression in plant cell culture has been described by Zenk, *Phytochemistry* 30, 3861-3863 (1991).

In particular, all plants from which protoplasts can be isolated and cultured to give whole regenerated plants can be utilised, so that whole plants are recovered which contain the transferred gene. Practically all plants can be regenerated from cultured cells or tissues, including but not limited to all major species of sugar cane, sugar beet, cotton, fruit and other trees, legumes and vegetables.

Examples of particularly preferred bacterial host cells include *streptococci*, *staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells.

Examples of particularly suitable host cells for fungal expression include yeast cells (for example, *S. cerevisiae*) and *Aspergillus* cells.

Any number of selection systems are known in the art that may be used to recover transformed cell lines. Examples include the herpes simplex virus thymidine kinase (Wigler, M. *et al.* (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. *et al.* (1980) *Cell* 22:817-23) genes that can be employed in  $tk^-$  or  $aprt^+$  cells, respectively.

Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dihydrofolate reductase (DHFR) that confers resistance to methotrexate (Wigler, M. *et al.* (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); *npt*, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin, F. *et al.* (1981) *J. Mol. Biol.* 150:1-14) and *als* or *pat*, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. Additional selectable genes have been

described, examples of which will be clear to those of skill in the art.

Although the presence or absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the relevant sequence is inserted within a marker gene sequence, transformed  
5 cells containing the appropriate sequences can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a polypeptide of the invention under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

- 10 Alternatively, host cells that contain a nucleic acid sequence encoding a polypeptide of the invention and which express said polypeptide may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassays, for example, fluorescence activated cell sorting (FACS) or immunoassay techniques (such as the enzyme-linked  
15 immunosorbent assay [ELISA] and radioimmunoassay [RIA]), that include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein (see Hampton, R. *et al.* (1990) Serological Methods, a Laboratory Manual, APS Press, St Paul, MN) and Maddox, D.E. *et al.* (1983) J. Exp. Med, 158, 1211-1216).

- A wide variety of labels and conjugation techniques are known by those skilled in the art  
20 and may be used in various nucleic acid and amino acid assays. Means for producing labelled hybridization or PCR probes for detecting sequences related to nucleic acid molecules encoding polypeptides of the present invention include oligolabelling, nick translation, end-labelling or PCR amplification using a labelled polynucleotide. Alternatively, the sequences encoding the polypeptide of the invention may be cloned into  
25 a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesise RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3 or SP6 and labelled nucleotides. These procedures may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, MI); Promega (Madison WI); and U.S. Biochemical Corp.,  
30 Cleveland, OH)).

Suitable reporter molecules or labels, which may be used for ease of detection, include radionuclides, enzymes and fluorescent, chemiluminescent or chromogenic agents as well

as substrates, cofactors, inhibitors, magnetic particles, and the like.

Nucleic acid molecules according to the present invention may also be used to create transgenic animals, particularly rodent animals. Such transgenic animals form a further aspect of the present invention. This may be done locally by modification of somatic cells, or by germ line therapy to incorporate heritable modifications. Such transgenic animals may be particularly useful in the generation of animal models for drug molecules effective as modulators of the polypeptides of the present invention.

The polypeptide can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulphate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. High performance liquid chromatography is particularly useful for purification. Well known techniques for refolding proteins may be employed to regenerate an active conformation when the polypeptide is denatured during isolation and or purification.

Specialised vector constructions may also be used to facilitate purification of proteins, as desired, by joining sequences encoding the polypeptides of the invention to a nucleotide sequence encoding a polypeptide domain that will facilitate purification of soluble proteins. Examples of such purification-facilitating domains include metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilised metals, protein A domains that allow purification on immobilised immunoglobulin, and the domain utilised in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, WA). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the polypeptide of the invention may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing the polypeptide of the invention fused to several histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilised metal ion affinity chromatography as described in Porath, J. *et al.* (1992), *Prot. Exp. Purif.* 3: 263-281) while the thioredoxin or enterokinase cleavage site provides a means for purifying the polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D.J. *et al.* (1993; *DNA Cell Biol.* 12:441-453).

If the polypeptide is to be expressed for use in screening assays, generally it is preferred that it be produced at the surface of the host cell in which it is expressed. In this event, the host cells may be harvested prior to use in the screening assay, for example using techniques such as fluorescence activated cell sorting (FACS) or immunoaffinity techniques. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the expressed polypeptide. If polypeptide is produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

The polypeptide of the invention can be used to screen libraries of compounds in any of a variety of drug screening techniques. Such compounds may activate (agonise) or inhibit (antagonise) the level of expression of the gene or the activity of the polypeptide of the invention and form a further aspect of the present invention. Preferred compounds are effective to alter the expression of a natural gene which encodes a polypeptide of the first aspect of the invention or to regulate the activity of a polypeptide of the first aspect of the invention.

Agonist or antagonist compounds may be isolated from, for example, cells, cell-free preparations, chemical libraries or natural product mixtures. These agonists or antagonists may be natural or modified substrates, ligands, enzymes, receptors or structural or functional mimetics. For a suitable review of such screening techniques, see Coligan *et al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991).

Compounds that are most likely to be good antagonists are molecules that bind to the polypeptide of the invention without inducing the biological effects of the polypeptide upon binding to it. Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to the polypeptide of the invention and thereby inhibit or extinguish its activity. In this fashion, binding of the polypeptide to normal cellular binding molecules may be inhibited, such that the normal biological activity of the polypeptide is prevented.

The polypeptide of the invention that is employed in such a screening technique may be free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. In general, such screening procedures may involve using appropriate cells or cell membranes that express the polypeptide that are contacted with a test compound to observe binding, or stimulation or inhibition of a functional response. The functional response of the cells contacted with the test compound is then compared with control cells that were

not contacted with the test compound. Such an assay may assess whether the test compound results in a signal generated by activation of the polypeptide, using an appropriate detection system. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist in the presence of the test compound is observed.

A preferred method for identifying an agonist or antagonist compound of a polypeptide of the present invention comprises:

(a) contacting a cell expressing on the surface thereof the polypeptide according to the first aspect of the invention, the polypeptide being associated with a second component capable of providing a detectable signal in response to the binding of a compound to the polypeptide, with a compound to be screened under conditions to permit binding to the polypeptide; and

(b) determining whether the compound binds to and activates or inhibits the polypeptide by measuring the level of a signal generated from the interaction of the compound with the polypeptide.

A further preferred method for identifying an agonist or antagonist of a polypeptide of the invention comprises:

(a) contacting a cell expressing on the surface thereof the polypeptide, the polypeptide being associated with a second component capable of providing a detectable signal in response to the binding of a compound to the polypeptide, with a compound to be screened under conditions to permit binding to the polypeptide; and

(b) determining whether the compound binds to and activates or inhibits the polypeptide by comparing the level of a signal generated from the interaction of the compound with the polypeptide with the level of a signal in the absence of the compound.

In further preferred embodiments, the general methods that are described above may further comprise conducting the identification of agonist or antagonist in the presence of labelled or unlabelled ligand for the polypeptide.

In another embodiment of the method for identifying agonist or antagonist of a polypeptide of the present invention comprises:

determining the inhibition of binding of a ligand to cells which have a polypeptide of the invention on the surface thereof, or to cell membranes containing such a polypeptide, in

the presence of a candidate compound under conditions to permit binding to the polypeptide, and determining the amount of ligand bound to the polypeptide. A compound capable of causing reduction of binding of a ligand is considered to be an agonist or antagonist. Preferably the ligand is labelled.

- 5 More particularly, a method of screening for a polypeptide antagonist or agonist compound comprises the steps of:
- (a) incubating a labelled ligand with a whole cell expressing a polypeptide according to the invention on the cell surface, or a cell membrane containing a polypeptide of the invention,
  - (b) measuring the amount of labelled ligand bound to the whole cell or the cell membrane;
  - 10 (c) adding a candidate compound to a mixture of labelled ligand and the whole cell or the cell membrane of step (a) and allowing the mixture to attain equilibrium;
  - (d) measuring the amount of labelled ligand bound to the whole cell or the cell membrane after step (c); and
  - (e) comparing the difference in the labelled ligand bound in step (b) and (d), such that the
  - 15 compound which causes the reduction in binding in step (d) is considered to be an agonist or antagonist.

The polypeptides may be found to modulate a variety of physiological and pathological processes in a dose-dependent manner in the above-described assays. Thus, the “functional equivalents” of the polypeptides of the invention include polypeptides that exhibit any of

20 the same modulatory activities in the above-described assays in a dose-dependent manner. Although the degree of dose-dependent activity need not be identical to that of the polypeptides of the invention, preferably the “functional equivalents” will exhibit substantially similar dose-dependence in a given activity assay compared to the polypeptides of the invention.

- 25 In certain of the embodiments described above, simple binding assays may be used, in which the adherence of a test compound to a surface bearing the polypeptide is detected by means of a label directly or indirectly associated with the test compound or in an assay involving competition with a labelled competitor. In another embodiment, competitive drug screening assays may be used, in which neutralising antibodies that are capable of
- 30 binding the polypeptide specifically compete with a test compound for binding. In this manner, the antibodies can be used to detect the presence of any test compound that

possesses specific binding affinity for the polypeptide.

Assays may also be designed to detect the effect of added test compounds on the production of mRNA encoding the polypeptide in cells. For example, an ELISA may be constructed that measures secreted or cell-associated levels of polypeptide using  
5 monoclonal or polyclonal antibodies by standard methods known in the art, and this can be used to search for compounds that may inhibit or enhance the production of the polypeptide from suitably manipulated cells or tissues. The formation of binding complexes between the polypeptide and the compound being tested may then be measured.

Assay methods that are also included within the terms of the present invention are those  
10 that involve the use of the genes and polypeptides of the invention in overexpression or ablation assays. Such assays involve the manipulation of levels of these genes/polypeptides in cells and assessment of the impact of this manipulation event on the physiology of the manipulated cells. For example, such experiments reveal details of signalling and metabolic pathways in which the particular genes/polypeptides are implicated, generate  
15 information regarding the identities of polypeptides with which the studied polypeptides interact and provide clues as to methods by which related genes and proteins are regulated.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the polypeptide of interest (see International patent application WO84/03564). In this method, large numbers of different  
20 small test compounds are synthesised on a solid substrate, which may then be reacted with the polypeptide of the invention and washed. One way of immobilising the polypeptide is to use non-neutralising antibodies. Bound polypeptide may then be detected using methods that are well known in the art. Purified polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques.

25 The polypeptide of the invention may be used to identify membrane-bound or soluble receptors, through standard receptor binding techniques that are known in the art, such as ligand binding and crosslinking assays in which the polypeptide is labelled with a radioactive isotope, is chemically modified, or is fused to a peptide sequence that facilitates its detection or purification, and incubated with a source of the putative receptor  
30 (for example, a composition of cells, cell membranes, cell supernatants, tissue extracts, or bodily fluids). The efficacy of binding may be measured using biophysical techniques such as surface plasmon resonance and spectroscopy. Binding assays may be used for the



purification and cloning of the receptor, but may also identify agonists and antagonists of the polypeptide, that compete with the binding of the polypeptide to its receptor. Standard methods for conducting screening assays are well understood in the art.

The invention also includes a screening kit useful in the methods for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, that are described above.

The invention includes the agonists, antagonists, ligands, receptors, substrates and enzymes, and other compounds which modulate the activity or antigenicity of the polypeptide of the invention discovered by the methods that are described above.

The invention also provides pharmaceutical compositions comprising a polypeptide, nucleic acid, ligand or compound of the invention in combination with a suitable pharmaceutical carrier. These compositions may be suitable as therapeutic or diagnostic reagents, as vaccines, or as other immunogenic compositions, as outlined in detail below.

According to the terminology used herein, a composition containing a polypeptide, nucleic acid, ligand or compound [X] is "substantially free of" impurities [herein, Y] when at least 85% by weight of the total X+Y in the composition is X. Preferably, X comprises at least about 90% by weight of the total of X+Y in the composition, more preferably at least about 95%, 98% or even 99% by weight.

The pharmaceutical compositions should preferably comprise a therapeutically effective amount of the polypeptide, nucleic acid molecule, ligand, or compound of the invention. The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent needed to treat, ameliorate, or prevent a targeted disease or condition, or to exhibit a detectable therapeutic or preventative effect. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, for example, of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

The precise effective amount for a human subject will depend upon the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. This amount can be determined by routine experimentation

and is within the judgement of the clinician. Generally, an effective dose will be from 0.01 mg/kg to 50 mg/kg, preferably 0.05 mg/kg to 10 mg/kg. Compositions may be administered individually to a patient or may be administered in combination with other agents, drugs or hormones.

- 5 A pharmaceutical composition may also contain a pharmaceutically acceptable carrier, for administration of a therapeutic agent. Such carriers include antibodies and other polypeptides, genes and other therapeutic agents such as liposomes, provided that the carrier does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity.
- 10 Suitable carriers may be large, slowly metabolised macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers and inactive virus particles.

Pharmaceutically acceptable salts can be used therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulphates, and the like; and the salts of  
15 organic acids such as acetates, propionates, malonates, benzoates, and the like. A thorough discussion of pharmaceutically acceptable carriers is available in Remington's Pharmaceutical Sciences (Mack Pub. Co., N.J. 1991).

Pharmaceutically acceptable carriers in therapeutic compositions may additionally contain liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such  
20 as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such compositions. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Once formulated, the compositions of the invention can be administered directly to the  
25 subject. The subjects to be treated can be animals; in particular, human subjects can be treated.

The pharmaceutical compositions utilised in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal or transcutaneous  
30 applications (for example, see WO98/20734), subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, intravaginal or rectal means. Gene guns or hyposprays may also be used to administer the pharmaceutical compositions of the invention. Typically, the

therapeutic compositions may be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared.

Direct delivery of the compositions will generally be accomplished by injection,  
5 subcutaneously, intraperitoneally, intravenously or intramuscularly, or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion. Dosage treatment may be a single dose schedule or a multiple dose schedule.

If the activity of the polypeptide of the invention is in excess in a particular disease state, several approaches are available. One approach comprises administering to a subject an  
10 inhibitor compound (antagonist) as described above, along with a pharmaceutically acceptable carrier in an amount effective to inhibit the function of the polypeptide, such as by blocking the binding of ligands, substrates, enzymes, receptors, or by inhibiting a second signal, and thereby alleviating the abnormal condition. Preferably, such antagonists are antibodies. Most preferably, such antibodies are chimeric and/or humanised to  
15 minimise their immunogenicity, as described previously.

In another approach, soluble forms of the polypeptide that retain binding affinity for the ligand, substrate, enzyme, receptor, in question, may be administered. Typically, the polypeptide may be administered in the form of fragments that retain the relevant portions.

In an alternative approach, expression of the gene encoding the polypeptide can be  
20 inhibited using expression blocking techniques, such as the use of antisense nucleic acid molecules (as described above), either internally generated or separately administered. Modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5' or regulatory regions (signal sequence, promoters, enhancers and introns) of the gene encoding the polypeptide.  
25 Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J.E. *et al.* (1994) In: Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches,  
30 Futura Publishing Co., Mt. Kisco, NY). The complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes. Such oligonucleotides may be administered or may be

generated *in situ* from expression *in vivo*.

In addition, expression of the polypeptide of the invention may be prevented by using ribozymes specific to its encoding mRNA sequence. Ribozymes are catalytically active RNAs that can be natural or synthetic (see for example Usman, N, *et al.*, Curr. Opin. Struct. Biol (1996) 6(4), 527-33). Synthetic ribozymes can be designed to specifically cleave mRNAs at selected positions thereby preventing translation of the mRNAs into functional polypeptide. Ribozymes may be synthesised with a natural ribose phosphate backbone and natural bases, as normally found in RNA molecules. Alternatively the ribozymes may be synthesised with non-natural backbones, for example, 2'-O-methyl RNA, to provide protection from ribonuclease degradation and may contain modified bases.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of non-traditional bases such as inosine, queosine and butosine, as well as acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine and uridine which are not as easily recognised by endogenous endonucleases.

For treating abnormal conditions related to an under-expression of the polypeptide of the invention and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound that activates the polypeptide, i.e., an agonist as described above, to alleviate the abnormal condition. Alternatively, a therapeutic amount of the polypeptide in combination with a suitable pharmaceutical carrier may be administered to restore the relevant physiological balance of polypeptide.

Gene therapy may be employed to effect the endogenous production of the polypeptide by the relevant cells in the subject. Gene therapy is used to treat permanently the inappropriate production of the polypeptide by replacing a defective gene with a corrected therapeutic gene.

Gene therapy of the present invention can occur *in vivo* or *ex vivo*. *Ex vivo* gene therapy requires the isolation and purification of patient cells, the introduction of a therapeutic

gene and introduction of the genetically altered cells back into the patient. In contrast, in vivo gene therapy does not require isolation and purification of a patient's cells.

The therapeutic gene is typically "packaged" for administration to a patient. Gene delivery vehicles may be non-viral, such as liposomes, or replication-deficient viruses, such as  
5 adenovirus as described by Berkner, K.L., in Curr. Top. Microbiol. Immunol., 158, 39-66 (1992) or adeno-associated virus (AAV) vectors as described by Muzyczka, N., in Curr. Top. Microbiol. Immunol., 158, 97-129 (1992) and U.S. Patent No. 5,252,479. For example, a nucleic acid molecule encoding a polypeptide of the invention may be engineered for expression in a replication-defective retroviral vector. This expression  
10 construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding the polypeptide, such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo (see Chapter 20, Gene Therapy and other Molecular  
15 Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics (1996), T Strachan and A P Read, BIOS Scientific Publishers Ltd).

Another approach is the administration of "naked DNA" in which the therapeutic gene is directly injected into the bloodstream or muscle tissue.

In situations in which the polypeptides or nucleic acid molecules of the invention are  
20 disease-causing agents, the invention provides that they can be used in vaccines to raise antibodies against the disease causing agent.

Vaccines according to the invention may either be prophylactic (ie. to prevent infection) or therapeutic (ie. to treat disease after infection). Such vaccines comprise immunising antigen(s), immunogen(s), polypeptide(s), protein(s) or nucleic acid, usually in  
25 combination with pharmaceutically-acceptable carriers as described above, which include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Additionally, these carriers may function as immunostimulating agents ("adjuvants"). Furthermore, the antigen or immunogen may be conjugated to a bacterial toxoid, such as a toxoid from diphtheria, tetanus, cholera, *H. pylori*, and other  
30 pathogens.

Since polypeptides may be broken down in the stomach, vaccines comprising polypeptides are preferably administered parenterally (for instance, subcutaneous, intramuscular,

intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient, and aqueous and non-aqueous sterile suspensions which may  
5 include suspending agents or thickening agents.

The vaccine formulations of the invention may be presented in unit-dose or multi-dose containers. For example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The dosage will depend on the specific activity of the vaccine and can be readily  
10 determined by routine experimentation.

Genetic delivery of antibodies that bind to polypeptides according to the invention may also be effected, for example, as described in International patent application WO98/55607.

The technology referred to as jet injection (see, for example, [www.powderject.com](http://www.powderject.com)) may  
15 also be useful in the formulation of vaccine compositions.

A number of suitable methods for vaccination and vaccine delivery systems are described in International patent application WO00/29428.

This invention also relates to the use of nucleic acid molecules according to the present invention as diagnostic reagents. Detection of a mutated form of the gene characterised by  
20 the nucleic acid molecules of the invention which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

25 Nucleic acid molecules for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR, ligase chain reaction (LCR), strand displacement amplification (SDA), or other amplification techniques (see Saiki *et al.*, Nature, 324, 163-166 (1986); Bej, *et al.*, Crit. Rev. Biochem.  
30 Molec. Biol., 26, 301-334 (1991); Birkenmeyer *et al.*, J. Virol. Meth., 35, 117-126 (1991); Van Brunt, J., Bio/Technology, 8, 291-294 (1990)) prior to analysis.

In one embodiment, this aspect of the invention provides a method of diagnosing a disease in a patient, comprising assessing the level of expression of a natural gene encoding a polypeptide according to the invention and comparing said level of expression to a control level, wherein a level that is different to said control level is indicative of disease. The method may comprise the steps of:

- a) contacting a sample of tissue from the patient with a nucleic acid probe under stringent conditions that allow the formation of a hybrid complex between a nucleic acid molecule of the invention and the probe;
  - b) contacting a control sample with said probe under the same conditions used in step a);
  - 10 c) and detecting the presence of hybrid complexes in said samples;
- wherein detection of levels of the hybrid complex in the patient sample that differ from levels of the hybrid complex in the control sample is indicative of disease.

A further aspect of the invention comprises a diagnostic method comprising the steps of:

- a) obtaining a tissue sample from a patient being tested for disease;
- 15 b) isolating a nucleic acid molecule according to the invention from said tissue sample; and
- c) diagnosing the patient for disease by detecting the presence of a mutation in the nucleic acid molecule which is associated with disease.

To aid the detection of nucleic acid molecules in the above-described methods, an amplification step, for example using PCR, may be included.

- 20 Deletions and insertions can be detected by a change in the size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labelled RNA of the invention or alternatively, labelled antisense DNA sequences of the invention. Perfectly-matched sequences can be distinguished from mismatched duplexes by RNase digestion or by assessing differences in melting
- 25 temperatures. The presence or absence of the mutation in the patient may be detected by contacting DNA with a nucleic acid probe that hybridises to the DNA under stringent conditions to form a hybrid double-stranded molecule, the hybrid double-stranded molecule having an unhybridised portion of the nucleic acid probe strand at any portion corresponding to a mutation associated with disease; and detecting the presence or absence
- 30 of an unhybridised portion of the probe strand as an indication of the presence or absence

of a disease-associated mutation in the corresponding portion of the DNA strand.

Such diagnostics are particularly useful for prenatal and even neonatal testing.

Point mutations and other sequence differences between the reference gene and "mutant" genes can be identified by other well-known techniques, such as direct DNA sequencing or single-strand conformational polymorphism, (see Orita *et al.*, Genomics, 5, 874-879 (1989)). For example, a sequencing primer may be used with double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures with radiolabelled nucleotides or by automatic sequencing procedures with fluorescent-tags. Cloned DNA segments may also be used as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with PCR. Further, point mutations and other sequence variations, such as polymorphisms, can be detected as described above, for example, through the use of allele-specific oligonucleotides for PCR amplification of sequences that differ by single nucleotides.

DNA sequence differences may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (for example, Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, Proc. Natl. Acad. Sci. USA (1985) 85: 4397-4401).

In addition to conventional gel electrophoresis and DNA sequencing, mutations such as microdeletions, aneuploidies, translocations, inversions, can also be detected by *in situ* analysis (see, for example, Keller *et al.*, DNA Probes, 2nd Ed., Stockton Press, New York, N.Y., USA (1993)), that is, DNA or RNA sequences in cells can be analysed for mutations without need for their isolation and/or immobilisation onto a membrane. Fluorescence *in situ* hybridization (FISH) is presently the most commonly applied method and numerous reviews of FISH have appeared (see, for example, Trachuck *et al.*, Science, 250, 559-562 (1990), and Trask *et al.*, Trends, Genet., 7, 149-154 (1991)).

In another embodiment of the invention, an array of oligonucleotide probes comprising a nucleic acid molecule according to the invention can be constructed to conduct efficient screening of genetic variants, mutations and polymorphisms. Array technology methods are well known and have general applicability and can be used to address a variety of



questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, Science (1996), Vol 274, pp 610-613).

In one embodiment, the array is prepared and used according to the methods described in PCT application WO95/11995 (Chee *et al.*); Lockhart, D. J. *et al.* (1996) Nat. Biotech. 14: 1675-1680); and Schena, M. *et al.* (1996) Proc. Natl. Acad. Sci. 93: 10614-10619).  
5 Oligonucleotide pairs may range from two to over one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support. In another aspect, an oligonucleotide may be synthesized on  
10 the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler *et al.*). In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such  
15 as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536 or 6144 oligonucleotides, or any other number between two and over one million which lends itself to the efficient use of commercially-available instrumentation.

20 In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (ant suitable solid support) and machines (including  
25 robotic instruments), and may contain 8, 24, 96, 384, 1536 or 6144 oligonucleotides, or any other number between two and over one million which lends itself to the efficient use of commercially-available instrumentation.

In addition to the methods discussed above, diseases may be diagnosed by methods comprising determining, from a sample derived from a subject, an abnormally decreased or  
30 increased level of polypeptide or mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for

instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

Assay techniques that can be used to determine levels of a polypeptide of the present invention in a sample derived from a host are well-known to those of skill in the art and are  
5 discussed in some detail above (including radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays). This aspect of the invention provides a diagnostic method which comprises the steps of: (a) contacting a ligand as described above with a biological sample under conditions suitable for the formation of a ligand-polypeptide complex; and (b) detecting said complex.

- 10 Protocols such as ELISA, RIA, and FACS for measuring polypeptide levels may additionally provide a basis for diagnosing altered or abnormal levels of polypeptide expression. Normal or standard values for polypeptide expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably humans, with antibody to the polypeptide under conditions suitable for complex formation  
15 The amount of standard complex formation may be quantified by various methods, such as by photometric means.

- Antibodies which specifically bind to a polypeptide of the invention may be used for the diagnosis of conditions or diseases characterised by expression of the polypeptide, or in assays to monitor patients being treated with the polypeptides, nucleic acid molecules,  
20 ligands and other compounds of the invention. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for the polypeptide include methods that utilise the antibody and a label to detect the polypeptide in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labelled by joining them, either covalently  
25 or non-covalently, with a reporter molecule. A wide variety of reporter molecules known in the art may be used, several of which are described above.

- Quantities of polypeptide expressed in subject, control and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease. Diagnostic assays may be used to  
30 distinguish between absence, presence, and excess expression of polypeptide and to monitor regulation of polypeptide levels during therapeutic intervention. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal

studies, in clinical trials or in monitoring the treatment of an individual patient.

A diagnostic kit of the present invention may comprise:

- (a) a nucleic acid molecule of the present invention;
- (b) a polypeptide of the present invention; or
- 5 (c) a ligand of the present invention.

In one aspect of the invention, a diagnostic kit may comprise a first container containing a nucleic acid probe that hybridises under stringent conditions with a nucleic acid molecule according to the invention; a second container containing primers useful for amplifying the nucleic acid molecule; and instructions for using the probe and primers for facilitating the  
10 diagnosis of disease. The kit may further comprise a third container holding an agent for digesting unhybridised RNA.

In an alternative aspect of the invention, a diagnostic kit may comprise an array of nucleic acid molecules, at least one of which may be a nucleic acid molecule according to the invention.

- 15 To detect polypeptide according to the invention, a diagnostic kit may comprise one or more antibodies that bind to a polypeptide according to the invention; and a reagent useful for the detection of a binding reaction between the antibody and the polypeptide.

Such kits will be of use in diagnosing a disease or disorder or susceptibility to disease or disorder in which alpha-2-macroglobulin-like proteinase inhibitors are implicated. Such  
20 diseases and disorders may include reproductive disorders, cell proliferative disorders, including neoplasm, melanoma, lung, colorectal, breast, pancreas, head and neck and other solid tumours; myeloproliferative disorders, such as leukemia, non-Hodgkin lymphoma, leukopenia, thrombocytopenia, angiogenesis disorder, Kaposi's sarcoma; autoimmune/inflammatory disorders, including allergy, inflammatory bowel disease,  
25 pancreatitis, arthritis, psoriasis, psoriasis vulgaris, respiratory tract inflammation, asthma, and organ transplant rejection; cardiovascular disorders, including hypertension, oedema, angina, atherosclerosis, thrombosis, sepsis, shock, reperfusion injury, and ischemia, particularly ischemic heart disease; neurological disorders including central nervous system disease, Alzheimer's disease, brain injury, Parkinson's disease, amyotrophic lateral  
30 sclerosis, and pain; developmental disorders; metabolic disorders including diabetes mellitus, osteoporosis, and obesity, AIDS, renal disease, particularly idiopathic nephrotic

syndrome; lung injury; infections including viral infection, bacterial infection, fungal infection and parasitic infection, particularly *Trypanosoma cruzi* infection and other pathological conditions. Preferably, the disease is one in which alpha-2-macroglobulin-like proteinase inhibitors are implicated.

- 5 Various aspects and embodiments of the present invention will now be described in more detail by way of example, with particular reference to the INSP087 and INSP088 polypeptides.

It will be appreciated that modification of detail may be made without departing from the scope of the invention.

#### 10 Brief description of the Figures

**Figure 1:** Top ten results from BLAST against NCBI non-redundant database using INSP087 polypeptide sequence.

**Figure 2:** Alignment generated by BLAST between the INSP087 polypeptide sequence and the closest annotated sequence, ovomacroglobulin ovastatin from *Gallus gallus*.

- 15 **Figure 3:** Top ten results from BLAST against NCBI non-redundant database using INSP088 polypeptide sequence.

**Figure 4:** Alignment generated by BLAST between the INSP088 polypeptide sequence and the closest annotated sequence, alpha-2-macroglobulin precursor from *Homo sapiens*.

- Figure 5:** INSP087 nucleotide sequence with translation showing predicted receptor  
20 binding domain (shaded).

**Figure 6:** Nucleotide (Figure 6a) and amino acid (Figure 6b) alignment of predicted INSP087 receptor binding domain and corresponding region of IMAGE clone 2243344.

**Figure 7:** INSP088 prediction nucleotide sequence with translation showing predicted receptor binding domain (shaded).

- 25 **Figure 8:** Nucleotide (Figure 8a) and amino acid (Figure 8b) sequence alignment of predicted INSP088 receptor binding domain and corresponding region of IMAGE clone 4753534.

**Figure 9:** Map of pENTR-INSP088-SP-6HIS

**Figure 10:** Map of pEAK12d-INSP088-SP-6HIS

**Figure 11:** Predicted nucleotide sequence of INSP088 with translation

**Figure 12:** Nucleotide sequence with translation of INSP088 PCR product cloned using primers INSP088-CP2 and INSP088-CP3

**Figure 13:** Map of pCR4-TOPO-INSP088-CP2/CP3

## 5 Examples

### Example 1: INSP087 protein BLAST results

The polypeptide sequence given in SEQ ID NO:68, which represents the translation of consecutive exons of INSP087, was used as a BLAST query against the NCBI non-redundant Sequence database.

- 10 The top ten matches are shown in Figure 1, all of which are alpha-2-macroglobulin-like proteinase inhibitors.

Figure 2 shows the alignment of the INSP087 query sequence to the sequence of the highest matching known protein, ovomacroglobulin ovastatin (*Gallus gallus*).

The INSP087 gene has been mapped to a chromosomal location 12p13.31

### 15 Example 2: INSP088 protein BLAST results

The polypeptide sequence given in SEQ ID NO:112, which represents the translation of consecutive exons of INSP088, was used as a BLAST query against the NCBI non-redundant Sequence database.

- 20 The top ten matches are shown in Figure 3, all of which are alpha-2-macroglobulin-like proteinase inhibitors.

Figure 4 shows the alignment of the INSP088 query sequence to the sequence of the highest matching known protein, alpha-2-macroglobulin precursor (*Homo sapiens*).

- Expressed sequence tags (ESTs) representing the INSP088 transcript in human originate from the following cDNA libraries: head-neck and brain. This suggests that INSP088 can  
25 be cloned from the above tissues and may be associated with diseases of the above tissues. Accordingly, the polypeptides, antibodies and other moieties described herein may have utility in the treating a disease in one of the above tissues.

The INSP088 gene has been mapped to a chromosomal location 12p13.31.

**Example 3: Cloning of receptor binding domain of INSP087**

A 95 amino acid region at the C-terminal end of the INSP087 alpha macroglobulin family member prediction was identified as the putative receptor binding domain of the INSP087 prediction.

- 5 Using this region to search public EST databases, several sequences were identified which matched the INSP087 receptor binding domain sequence. One sequence was chosen, GenBank Accession AI655905 corresponding to IMAGE cDNA clone 2243344, and the clone was purchased from ATCC. The insert sequence of the IMAGE clone was sequenced using sequencing primers T7 and T3 (Table 1). The insert sequence was found to contain  
10 the predicted INSP087 receptor binding domain. IMAGE clone 2243344 is plasmid database ID 13434.

The sequence of the INSP087 prediction with position of the receptor binding domain is shown in Figure 5. The alignment of the nucleic acid and amino acid sequences of the INSP087 receptor binding domain with the corresponding region of IMAGE clone  
15 2243344 are shown in Figures 6a and 6b respectively.

Table 1: IMAGE clone 2243344 sequencing primers:

Primer	Sequence (5'-3')
T7 primer	TAA TAC GAC TCA CTA TAG G
T3 primer	ATT AAC CCT CAC TAA AGG

**Example 4 : Cloning of receptor binding domain of INSP088**

20 **1. Cloning of INSP088 receptor binding domain**

- A 92 amino acid region at the C-terminal end of the INSP088 alpha macroglobulin family member prediction was identified as encoding the receptor binding domain. The sequence of the INSP088 prediction with the receptor binding domain highlighted is shown in Figure 7. Using this domain to search public EST databases, we identified several sequences  
25 which matched the INSP088 receptor binding domain sequence. IMAGE cDNA clone 4753534, (GenBank Accession BG680886) was subsequently purchased from ATCC.

The insert sequence of the IMAGE clone was sequenced using sequencing primers T7 and SP6 (Table 2). The insert sequence was found to contain the predicted INSP088 receptor binding domain. The alignment of the nucleotide and amino acid sequences of the

INSP088 receptor binding domain with the corresponding region of IMAGE clone 4753534 is shown in Figures 8a and 8b respectively. IMAGE clone 4753534 is plasmid database ID 13684.

2. Subcloning of the INSP088 ORF with a C-terminal 6HIS tag and an IL12p40  
5 signal sequence into pEAK12d for expression in HEK293/EBNA cells

The INSP088 ORF fused to a C-terminal 6HIS tag and a N-terminal signal peptide sequence from IL12p40 was subcloned into pEAK12d mammalian cell using the Gateway cloning methodology.

The first stage of the Gateway cloning process involves a 3 step PCR reaction which  
10 generates the ORF of INSP088 flanked at the 5' end by an attB1 recombination site, a Kozak sequence and the IL12p40 signal sequence, and flanked at the 3' end by a sequence encoding an in-frame 6 histidine (6HIS) tag, a stop codon and the attB2 recombination site (Gateway compatible cDNA).

The first PCR reaction (in a final volume of 50 µl) contains: 1.5 µl of miniprep DNA  
15 prepared from IMAGE clone 4753534 ( plasmid ID 13684), 1.5 µl dNTPs (10 mM), 5 µl of 10X Pfx polymerase buffer, 1 µl MgSO<sub>4</sub> (50 mM), 0.5 µl each of gene specific primer (100 µM) (INSP088-PCR1F and INSP088-PCR1R) and 0.5 µl Platinum Pfx DNA polymerase (Invitrogen). The PCR reaction was performed using an initial denaturing step of 95°C for 1 min, followed by 10 cycles of 94 °C for 15 s; 55 °C for 30 s and 68°C for 1  
20 min 30 s; and a holding cycle of 4 °C. PCR products were purified directly from the reaction mixture using the Wizard PCR prep DNA purification system (Promega) according to the manufacturer's instructions.

The second PCR reaction (in a final volume of 50 µl) contained 10 µl purified PCR product, 1.5 µl dNTPs (10 mM), 5 µl of 10X Pfx polymerase buffer, 1 µl MgSO<sub>4</sub> (50  
25 mM), 0.5 µl of IL12p40 signal sequence forward primer, IL12P40-SP1 (100 µM), 0.5 µl GCP reverse primer (100 µM) and 0.5 µl of Platinum Pfx DNA polymerase. The conditions for the 2nd PCR reaction were: 95 °C for 1 min, followed by 10 cycles of 94 °C for 15 s; 55 °C for 30 s and 68°C for 1 min 30 s; and a holding cycle of 4 °C. PCR products were purified directly from the reaction mixture using the Wizard PCR prep DNA  
30 purification system (Promega) according to the manufacturer's instructions.

The third PCR reaction (in a final volume of 50 µl) contained 10 µl purified PCR product, 1.5 µl dNTPs (10 mM), 5 µl of 10X Pfx polymerase buffer, 1 µl MgSO<sub>4</sub> (50 mM), 0.5 µl of Gateway conversion signal peptide forward primer, GCPF-SP (100 µM), 0.5 µl GCP reverse primer (100 µM) and 0.5 µl of Platinum Pfx DNA polymerase. The conditions for the third PCR reaction were: 95 °C for 1 min; 4 cycles of 94 °C, 15 sec; 50 °C, 30 sec and 68 °C for 1 min 30 s; 25 cycles of 94 °C, 15 sec; 55 °C, 30 sec and 68 °C, 1 min 30 s; followed by a holding cycle of 4 °C. PCR products were gel purified using the Wizard PCR prep DNA purification system and eluted in 50 µl of sterile water.

10        3. Subcloning of Gateway compatible INSP088 ORF into Gateway entry vector pDONR201 and expression vector pEAK12d

The second stage of the Gateway cloning process involves subcloning of the Gateway modified PCR product into the Gateway entry vector pDONR201 as follows: 5 µl of purified PCR product is incubated with 1.5 µl pDONR201 vector (0.1 µg/µl), 2 µl BP buffer and 1.5 µl of BP clonase enzyme mix (Invitrogen) at RT for 2 h. The reaction was stopped by addition of proteinase K (2 µg) and incubated at 37 °C for a further 10 min.

An aliquot of this reaction (1 µl) was used to transform E. coli DH10B cells by electroporation as follows: a 20 µl aliquot of DH10B electrocompetent cells (Invitrogen) was thawed on ice and 1 µl of the BP reaction mix was added. The mixture was transferred to a chilled 0.1 cm electroporation cuvette and the cells electroporated using a BioRad Gene-Pulser™ according to the manufacturer's recommended protocol. SOC media (1 ml) which had been pre-warmed to room temperature was added immediately after electroporation. The mixture was transferred to a 15 ml snap-cap tube and incubated, with shaking (220 rpm) for 1 h at 37 °C. Aliquots of the transformation mixture (100 µl) were then plated on L-broth (LB) plates containing kanamycin (40 µg/ml) and incubated overnight at 37 °C. Transformants were plated on LB-kanamycin plates.

Plasmid mini-prep DNA was prepared from 5 ml cultures from 6 of the resultant colonies using a Qiaprep Turbo 9600 robotic system (Qiagen). Plasmid DNA (200-500 ng) was subjected to DNA sequencing with 21M13 and M13Rev primers using the BigDyeTerminator system (Applied Biosystems cat. no. 4390246) according to the manufacturer's instructions. The primer sequences are shown in Table 2. Sequencing reactions were purified using Dye-Ex columns (Qiagen) or Montage SEQ 96 cleanup



plates (Millipore cat. no. LSKS09624) then analyzed on an Applied Biosystems 3700 sequencer.

Sequence verified miniprep DNA (pENTR-INSP088-SP-6HIS, plasmid ID 13424, figure 9) 3 µl, was then used in a recombination reaction containing 1.5 µl pEAK12d vector (0.1 µg / µl), 2 µl LR buffer and 1.5 µl of LR clonase (Invitrogen) in a final volume of 10 µl. The mixture was incubated at RT for 1 h, stopped by addition of proteinase K (2 µg) and incubated at 37°C for a further 10 min. An aliquot of this reaction (1 µl) was used to transform *E. coli* DH10B cells by electroporation as follows: a 20 µl aliquot of DH10B electrocompetent cells (Invitrogen) was thawed on ice and 1 µl of the BP reaction mix was added. The mixture was transferred to a chilled 0.1 cm electroporation cuvette and the cells electroporated using a BioRad Gene-Pulser™ according to the manufacturer's recommended protocol. SOC media (1 ml) which had been pre-warmed to room temperature was added immediately after electroporation. The mixture was transferred to a 15 ml snap-cap tube and incubated, with shaking (220 rpm) for 1 h at 37 °C. Aliquots of the transformation mixture (100 µl) were then plated on L-broth (LB) plates containing ampicillin (100 µg/ml) and incubated overnight at 37 °C.

Plasmid mini-prep DNA was prepared from 5 ml cultures from 4 of the resultant colonies using a Qiaprep Turbo 9600 robotic system (Qiagen). Plasmid DNA (200-500 ng) was subjected to DNA sequencing with pEAK12d F and pEAK12d R primers as described above.

CsCl gradient purified maxi-prep DNA of plasmid pEAK12d-INSP088- SP-6HIS (plasmid ID number 13439) (figure 10) was prepared from a 500 ml culture of sequence verified clones (Sambrook J. et al., in Molecular Cloning, Laboratory Manual, 2<sup>nd</sup> edition, 1989, Cold Spring Harbor Laboratory Press), resuspended at a concentration of 1 µg/µl in sterile water and stored at -20 C.

Table 2: INSP088 cloning and sequencing primers

Primer	Sequence (5' -3')
INSP088-PCR1F	TTT TCC CTG GTT TTT CTG GCA TCT CCC CTC GTG GCC ATT GTG GAA GTG AA ATG CT
INSP088-PCR1R	GTG ATG GTG ATG GTG ACA GGG ATC AGA ATA CTG AA

IL12P40- SP1	<u>ATG TGT CAC CAG CAG TTG GTC ATC TCT TGG TTT TCC CTG GTT TTT CTG GCA</u> <u>TCT CCC CTC GTG GCC ATA</u>
GCPF-SP	GGGG ACA AGT TTG TAC AAA AAA GCA GGC TTC <u>GCC ACC</u> <u>ATG TGT CAC CAG CAG</u> <u>TTG</u>
GCP Reverse	GGG GAC CAC TTT GTA CAA GAA AGC TGG GTT TCA <i>ATG GTG ATG GTG ATG GTG</i>
T7 primer	TAA TAC GAC TCA CTA TAG G
SP6 primer	ATT TAG GTG ACA CTA TAG
pEAK12F	GCC AGC TTG GCA CTT GAT GT
pEAK12R	GAT GGA GGT GGA CGT GTC AG
21M13	TGT AAA ACG ACG GCC AGT
M13REV	CAG GAA ACA GCT ATG ACC

Underlined sequence = Kozak sequence

**Bold** = Stop codon

*Italic* sequence = His tag

5 **Bold underlined** = IL12p40 signal sequence

### **Example 5 : Cloning of macroglobulin domain of INSP088**

#### **1. Preparation of human cDNA templates**

- First strand cDNA was prepared from a variety of normal human tissue total RNA samples (purchased from Clontech, Stratagene, Ambion, Biochain Institute or prepared in-house)
- 10 using Superscript II RNase H<sup>-</sup> Reverse Transcriptase (Invitrogen) according to the manufacturer's protocol. Oligo (dT)<sub>15</sub> primer 1 µl (500 µg/ml, Promega), 2 µg human total RNA, 1 µl 10 mM dNTP Mix (10 mM each of dATP, dGTP, dCTP and dTTP at neutral pH) and sterile distilled water to a final volume of 12 µl were combined in a 1.5 ml Eppendorf tube, heated to 65 °C for 5 min and then chilled on ice.
- 15 The contents of the tube were collected by brief centrifugation and 4 µl 5X First-Strand Buffer, 2 µl of 0.1 M DTT, and 1 µl of RnaseOUT Recombinant Ribonuclease Inhibitor (40 units/µl, Invitrogen) were added. The contents of the tube were mixed gently and incubated at 42 °C for 2 min, then 1 µl (200 units) of SuperScript II enzyme was added and mixed gently by pipetting. The mixture was incubated at 42 °C for 50 min and then
- 20 inactivated by heating at 70 °C for 15 min. To remove RNA complementary to the cDNA, 1 µl (2 units) of *E. coli* RNase H (Invitrogen) was added and the reaction mixture incubated at 37 °C for 20 min. The final 21 µl reaction mix was diluted by adding 179 µl

sterile water to give a total volume of 200 µl. One µl of each cDNA sample was used as a template for PCR in a 50 µl final reaction volume, which corresponded to approximately 20 ng of cDNA.

## 2. Gene specific cloning primers for PCR

- 5 A pair of PCR primers having a length of between 18 and 25 bases were designed for amplifying the predicted macroglobulin domain of the INSP088 prediction. Primers were designed using Primer Designer Software (Scientific & Educational Software, PO Box 72045, Durham, NC 27722-2045, USA). PCR primers were optimized to have a  $T_m$  close to  $55 \pm 10$  °C and a GC content of 40-60%. Primers were selected which had high  
10 selectivity for the target sequence (little or no non specific priming).

## 3. PCR amplification of INSP088 macroglobulin domain from human cDNA

Gene-specific cloning primers INSP088-CP2 and INSP088-CP3 (Figures 11 & 12 and Table 3) were designed to amplify a cDNA fragment of 2252 bp covering 446 – 2697 bp of the coding sequence of the INSP088 prediction.

- 15 The primer pair was used with human cDNA samples from brain, testis, colon and skin as templates. The PCR was performed in a final volume of 50 µl containing 1X Easy-A™ reaction buffer, 200 µM dNTPs, 15 pmoles each of cloning primer, 2.5 units of Easy-A™ PCR polymerase (Stratagene) and approximately 20 ng of each cDNA template using an MJ Research DNA Engine, programmed as follows: 95 °C, 2 min; 40 cycles of 95 °C, 40  
20 sec, 53 °C, 30 sec, and 72 °C, 3 min; followed by 1 cycle at 72 °C for 7 min and a holding cycle at 4 °C. The amplification products were visualized on 0.8 % agarose gels in 1 X TAE buffer (Invitrogen). PCR products migrating at the predicted molecular mass were purified from the gel using the MinElute Gel Extraction System (QIAGEN). The PCR products were eluted in 10 µl of buffer EB (10mM Tris.Cl, pH8.5) and subcloned directly.

## 4. Subcloning of PCR Products

25 PCR products were subcloned into the topoisomerase I modified cloning vector (pCR4-TOPO) using the TOPO cloning kit purchased from the Invitrogen Corporation using the conditions specified by the manufacturer.

- Briefly, 4 µl of gel purified PCR product from the human cDNA amplification was  
30 incubated for 15 min at room temperature with 1 µl of TOPO vector and 1 µl salt solution. The reaction mixture was then transformed into *E. coli* strain TOP10 (Invitrogen) as follows: a 50 µl aliquot of One Shot TOP10 cells was thawed on ice and 2 µl of TOPO

reaction was added. The mixture was incubated for 15 min on ice and then heat shocked by incubation at 42 °C for exactly 30 s. Samples were returned to ice and 250 µl of warm SOC media (room temperature) was added. Samples were incubated with shaking (220 rpm) for 1 h at 37 °C. The transformation mixture was then plated on L-broth (LB) plates containing  
5 ampicillin (100 µg/ml) and incubated overnight at 37 °C. Ampicillin resistant colonies containing inserts were identified by colony PCR.

### 5. Colony PCR

Colonies were inoculated into 50 µl sterile water using a sterile toothpick. A 10 µl aliquot  
10 of the inoculum was then subjected to PCR using amplification primers INSP088-CP1 and INSP088-CP2 (Table 3). This pair of primers had been designed to amplify a 307 bp product containing the receptor binding domain of INSP088 located at the 3' end of the macroglobulin domain. This receptor binding domain had previously been identified in IMAGE clone 4753534 (plasmid ID 13443) which was used as a positive control template  
15 for the colony PCR. The PCR was performed in a final volume of 20 µl containing 1X AmpliTaq™ buffer, 200 µM dNTPs, 20 pmoles each of cloning primer, 1 unit of AmpliTaq™ (Perkin Elmer) and 10 µl of the bacterial colony solution using an MJ Research DNA Engine, programmed as follows: 94 °C, 2 min; 30 cycles of 94 °C, 30 sec, 55 °C, 30 sec and 72 °C for 30 sec. Samples were then maintained at 4 °C (holding cycle)  
20 before further analysis.

PCR products were analyzed on 1% agarose gels in 1 X TAE buffer. Colonies which gave approximately the expected PCR product size (307 bp + 187 bp due to the multiple cloning site or MCS) were grown up overnight at 37 °C in 5 ml L-Broth (LB) containing ampicillin (100µg /ml), with shaking at 220 rpm.

### 25 6. Plasmid DNA preparation and sequencing

Miniprep plasmid DNA was prepared from 5 ml cultures using a Qiaprep Turbo 9600 robotic system (Qiagen) or Wizard Plus SV Minipreps kit (Promega cat. no. 1460) according to the manufacturer's instructions. Plasmid DNA was eluted in 100 µl of sterile water. The DNA concentration was measured using an Eppendorf BO photometer. Plasmid  
30 DNA (200-500 ng) was subjected to DNA sequencing with the T7 primer and T3 primer using the BigDyeTerminator system (Applied Biosystems cat. no. 4390246) according to the manufacturer's instructions. The primer sequences are shown in Table 3. Sequencing reactions were purified using Dye-Ex columns (Qiagen) or Montage SEQ 96 cleanup

plates (Millipore cat. no. LSKS09624) then analyzed on an Applied Biosystems 3700 sequencer. Clones which were identified as containing the expected insert after sequencing with the T7 and T3 primers were subjected to further sequencing with gene-specific sequencing primers INSP088-SP1, INSP088-SP2 and INSP088-SP3 (Table 3).

- 5 Sequence analysis identified a clone which contained 100% match to the predicted INSP088-CP2/INSP088-CP3 product sequence. The sequence of this cloned cDNA fragment is shown in Figure 12. The plasmid map of the cloned PCR product pCR4-TOPO-INSP088-CP2/CP3 (plasmid ID 13684) is shown in Figure 13.

10 Table 3: INSP088 cloning and sequencing primers

Primer	Sequence (5'-3')
INSP088-CP1	GAG CCG TAG CTC TTC CAA TA
INSP088-CP2	GCT GGA CTT ACC TCA TTC AC
INSP088-CP3	GTG GTC ATC CAG AGG CTT T
INSP088-SP1	GGA CAG CAA TGA ACC ATG TG
INSP088-SP2	CAC AGC AAT GGC TCA TAC AG
INSP088-SP3	AGC CAG AAA CCT ACT CCA TC
T7 primer	TAA TAC GAC TCA CTA TAG G
T3 primer	ATT AAC CCT CAC TAA AGG

#### **Example 6 : Assays for determining biological activity**

##### **Differentiation to adipocyte assay:**

- Inhibition of adipocyte differentiation is an in vitro model for reduction of adipose mass  
 15 believed to be important in reducing insulin resistance in diseases such as diabetes and Polycystic Ovary Syndrome (PCOS). The goal is to identify protein(s) that inhibits differentiation of pre-adipocytes to adipocytes. The 3T3-L1 mouse preadipocyte cell line is induced to differentiate to adipocytes with insulin + IBMX. That differentiation is inhibited by TNF-alpha + cyclohexamide as a positive control.

20

##### **Tritiated glucose uptake (3T3 L1):**

The goal of this assay is to identify protein (s) that stimulate glucose uptake as a model for insulin-resistance in adipose during diabetes or PCOS. Adipocytes used are mouse 3T3-L1 preadipocytes that have been differentiated.

· Tritiated glucose uptake (primary human adipocytes):

The goal of this assay is to identify protein (s) that stimulate glucose uptake as a model for insulin-resistance in adipose during diabetes or PCOS. Primary human adipocytes are used.

5

· Tritiated glucose uptake (primary human skeletal muscle cells)

The goal of this assay is to identify protein (s) that stimulate glucose uptake as a model for Insulin-resistance in muscle tissue during diabetes or PCOS. Primary human skeletal muscle cells are differentiated into myotubes and then used in the assay.

10

**INSP087 and INSP088 sequences (Note: for amino acids encoded by exon-exon junctions, the amino acid will be assigned to the more 5' exon.)**

**SEQ ID NO: 1 (INSP087 nucleotide sequence exon 1)**

5       1 GTTCCTCAGG CCAGATCTGA CCCACTGGCA TTTATTACAT TTTCTGCTAA  
      51 AGGAGCCACT CTCAACCTGG AAGAGAGGAG ATCTGTGGCA ATCAGATCCA  
      101 GAGAGAATGT GGTCTTCGTA CAGACTGATA AACCCACCTA CAAGCCTGGA  
      151 CAGAAAG

10 **SEQ ID NO: 2 (INSP087 protein sequence exon 1)**

      1 VPQARSDPLA FITFSAKGAT LNLEERRSVA IRSRENVVFV QTDKPTYKPG  
      51 QKV

**SEQ ID NO: 3 (INSP087 nucleotide sequence exon 2)**

15       1 TTCATATATT AACATTATTT TTATTTTAT TTCAGTATCC AGTGATCACC  
      51 CTTCAG

**SEQ ID NO: 4 (INSP087 protein sequence exon 2)**

      1 HILTLFLFLF QYPVITLQ

20

**SEQ ID NO: 5 (INSP087 nucleotide sequence exon 3)**

      1 GATCCTCAAA ACAATCGGAT TTTTCAAAGG CAAAATGTGA CTTCTTTCCG  
      51 AAATATTACC CAACTCTCGT TCCAACTGAT TTCAGAACCA ATGTTTGGAG  
      101 ATTACTGGAT TGTGTGAAA AGAAACTCAA GGGAGACAGT GACACACCAA  
25       151 TTGCTGTTA AAAGATATG

**SEQ ID NO: 6 (INSP087 protein sequence exon 3)**

      1 DPQNNRIFQR QNVTSFRNIT QLSFQLISEP MFGDYWIVVK RNSRETVTHQ  
      51 FAVKRYV

30

**SEQ ID NO: 7 (INSP087 nucleotide sequence exon 4)**

      1 TGCTGCCCAA GTTTGAAGTT ACAGTCAATG CACCACAAAC AGTAACTATT  
      51 TCAGATGATG AATTCCAAGT GGATGTATGT GCTAA

**SEQ ID NO: 8 (INSP087 protein sequence exon 4)**

1 LPKFEVTVNA PQTVTISDDE FQVDVCAK

**5 SEQ ID NO: 9 (INSP087 nucleotide sequence exon 5)**

1 GTACAACTTT GGCCAACCTG TGCAAGGGGA AACCCAAATC CGGGTGTGCA  
51 GAGAGTATTT TTCTTCAAGC AATTGTGAGA AAAATGAAAA TGAAATATGT  
101 GAGCAATTTA TTGCACAG

**10 SEQ ID NO: 10 (INSP087 protein sequence exon 5)**

1 YNFGQPVQGE TQIRVCREYF SSSNCEKNEN EICEQFIAQ

**SEQ ID NO: 11 (INSP087 nucleotide sequence exon 6)**

1 TTGGAAAATG GTTGTGTTTC TCAAATTGTA AATACAAAAG TCTTCCAAC  
15 51 CTACCGTTCG GGATTGTTCA TGACATTTCA TGTCGCTGTA ATTGTTACAG  
101 AATCTGGGAC AG

**SEQ ID NO: 12 (INSP087 protein sequence exon 6)**

1 LENGCVSQIV NTKVFQLYRS GLFMTFHVAV IVTESGTV

20

**SEQ ID NO: 13 (INSP087 nucleotide sequence exon 7)**

1 TTATGCAGAT CAGCGAGAAG ACCTCAGTTT TTATCACTCA ATTGCTTGGA  
51 ACTGTAAACT TTGAGAACAT GGATACATTC TATAGAAGAG GGATTTCTTA  
101 TTTTGGAAC

25

**SEQ ID NO: 14 (INSP087 protein sequence exon 7)**

1 MQISEKTSVF ITQLLGTVNF ENMDTFYRRG ISYFGT

**SEQ ID NO: 15 (INSP087 nucleotide sequence exon 8)**

30 1 CTTAAATTTT CGGATCCCAA TAATGTACCT ATGGTGAACA AGTTGTTGCA  
51 ACTGGAGCTC AATGATGAAT TTATAGGAAA TTACACTACG GATGAGAATG  
101 GCGAAGCTCA ATTTTCCATT GACACTTCAG ACATATTGA TCCAGAGTTC  
151 AACCTAAAA



**SEQ ID NO: 16 (INSP087 protein sequence exon 8)**

1 LKFSDPNNVP MVNKLLQLEL NDEFIGNYTT DENGEAQFSI DTSDIFDPEF  
51 NLK

5

**SEQ ID NO: 17 (INSP087 nucleotide sequence exon 9)**

1 GCCACATATG TTCGACCTGA GAGCTGCTAT CTTCCCAGCT GGTGACGCC  
51 TCAGTACTTG GATGCTCACT TCTTAGTCTC ACGCTTTTAC TCCCGAACCA  
101 ACAGCTTCCT GAAGATTGTT CCAGAACCAA AGCAGCTTGA ATGTAATCAA  
10 151 CAGAAGGTTG TTAGTGTGCA TTAGTCCCTA AACAGTGAAG CATATGAGGA  
201 TGATTCCAAT GTAAAGTTCT TCTATTTG

**SEQ ID NO: 18 (INSP087 protein sequence exon 9)**

1 ATYVRPESCY LPSWLTPQYL DAHFLVSRFY SRTNSFLKIV PEPKQLECNQ  
15 51 QKVVTVHYSL NSEAYEDDSN VKFFYL

**SEQ ID NO: 19 (INSP087 nucleotide sequence exon 10)**

1 ATGATGGTAA AAGGAGCTAT CTTACTCAGT GGACAAAAGG AAATCAGAAA  
51 CAAAG  
20

**SEQ ID NO: 20 (INSP087 protein sequence exon 10)**

1 MMVKGAILLS GQKEIRNKA

**SEQ ID NO: 21 (INSP087 nucleotide sequence exon 11)**

25 1 CCTGGAATGG AACTTCTCG TTCCCAATCA GCATCAGTGC TGATCTGGCT  
51 CCTGCAGCCG TCCTGTTTGT CTATACCCTT CACCCCAGTG GGGAAATTGT  
101 GGCTGACAGT GTCAGATTCC AGGTTGACAA GTGCTTTAAA CACAAG

**SEQ ID NO: 22 (INSP087 protein sequence exon 11)**

30 1 WNGNFSFPIS ISADLAPAAV LFVYTLHPSG EIVADSVRFQ VDKCFKHK

**SEQ ID NO: 23 (INSP087 nucleotide sequence exon 12)**

1 GTTAACATAA AGTTCTCTAA CGAGCAGGGC TTACCTGGTT CCAATGCTAG  
51 TCTCTGTCTT CAAGCGGCGC CTGTCTTATT CTGTGCCCTC AGGGCTGTGG

101 ATAGGAATGT CCTTCTACTG AAATCTGAAC AACAGCTGTC AGCTGAAAGT

**SEQ ID NO: 24 (INSP087 protein sequence exon 12)**

1 VNIKFSNEQG LPGSNASLCL QAAPVLFCAL RAVDRNVLLL KSEQQLSAES

5

**SEQ ID NO: 25 (INSP087 nucleotide sequence exon 13)**

1 GTGTATAACA TGGTTCCAAG TATAGAGCCG TATGGTTATT TCTACCATGG

51 CCTCAATCTT GATGATGGCA AGGAAGACCC TTGCATTCCT CAGAGGGATA

101 TGTTCTACAA TGGTTTATAT TACACACCTG TAAGCAACTA TGGGGATGGA

10 151 GATATCTATA ATATTGTCAG G

**SEQ ID NO: 26 (INSP087 protein sequence exon 13)**

1 VYNMVPSIEP YGYFYHGLNL DDGKEDPCIP QRDMFYNGLY YTPVSNYGDG

51 DIYNIVR

15

**SEQ ID NO: 27 (INSP087 nucleotide sequence exon 14)**

1 AACATGGGTC TAAAAGTCTT TACCAATCTC CATTACCGAA AACCAGAAGT

51 ATGTGTGATG GAGAGAAGGC TGCCACTCCC TAAGCCGCTT TATCTGGAAA

101 CAGAAAATTA TGGTCCAATG CGTAGTGTTT CGTCTAGAAT TGCATGTAG

20

**SEQ ID NO: 28 (INSP087 protein sequence exon 14)**

1 NMGLKVFTNL HYRKPEVCVM ERRLPLPKPL YLETENYGPM RSVPSRIACR

**SEQ ID NO: 29 (INSP087 nucleotide sequence exon 15)**

25 1 AGGGGAGAAT GCTGACTATG TAGAACAGGC TATAATTCAA ACAGTAAGAA

51 CAAACTTCCC AGAGACATGG ATGTGGGACC TCGTCAGTGT CGA

**SEQ ID NO: 30 (INSP087 protein sequence exon 15)**

1 GENADYVEQA IIQTVRTNFP ETWMWDLVSV D

30

**SEQ ID NO: 31 (INSP087 nucleotide sequence exon 16)**

1 TTCCTCAGGC TCTGCCAATC TTTCGTTTCTT CATTCCTGAT ACGATAACCC

51 AATGGGAGGC AAGTGGCTTT TGTGTGAATG GTGACGTTGG ATTTGGCATT

101 TCCTCTACAA CCACTCTAGA AGTCTCCCAA CCTTTCTTTA TTGAGATTGC

151 CTCACCCTTT TCGGTTGTTT AAAATGAACA ATTTGATTTG ATTGTCAATG  
201 TCTTCAGCTA CCGGAATACA TGTGTAGAG

**SEQ ID NO: 32 (INSP087 protein sequence exon 16)**

5 1 SSGSANLSFL IPDTITQWEA SGFCVNGDVG FGISSTTLE VSQPFIEIA  
51 SPFSVVQNEQ FDLIVNVFSY RNTCVE

**SEQ ID NO: 33 (INSP087 nucleotide sequence exon 17)**

1 ATTTCTGTTC AAGTGGAGGA GTCTCAGAAT TATGAAGCAA ATATTCATAC  
10 51 CTTGAAAATC AATGGCAGTG AGGTTATTCA AGCTGGAGGG AGGAAAACAA  
101 ACGTCTGGAC TATTATACCT AAGAAATTGG

**SEQ ID NO: 34 (INSP087 protein sequence exon 17)**

1 ISVQVEESQN YEANIHTLKI NGSEVIQAGG RKTNVWTIIP KKLK  
15

**SEQ ID NO: 35 (INSP087 nucleotide sequence exon 18)**

1 GTAAAGTGAA TATCACTGTA GTTGCTGAGT CCAAACAAAG CAGTGCTTGC  
51 CCAAATGAAG GAATGGAGCA GCAAAAGCTA AACTGGAAAG AACTGTGGT  
101 CCAAAGCTTC TTAGTAGAG  
20

**SEQ ID NO: 36 (INSP087 protein sequence exon 18)**

1 KVNITVVAES KQSSACPNEG MEQQKLNWKD TVVQSFLVE

**SEQ ID NO: 37 (INSP087 nucleotide sequence exon 19)**

25 1 CCTGAAGGTA TTGAAAAGGA AAGGACCCAG AGTTTCCTTA TCTGTACAGA  
51 AG

**SEQ ID NO: 38 (INSP087 protein sequence exon 19)**

1 PEGIEKERTQ SFRICTEG  
30

**SEQ ID NO: 39 (INSP087 nucleotide sequence exon 20)**

1 GTGCCAAAGC CTCCAAGCAG GGAGTTTGG ACTTGCCAAA CGATGTAGTA  
51 GAAGGGTCAG CCAGAGGCTT TTCTACTGTT GTGG

**SEQ ID NO: 40 (INSP087 protein sequence exon 20)**

1 AKASKQGVLD LPNDVVEGSA RGFFTIVVG

**5 SEQ ID NO: 41 (INSP087 nucleotide sequence exon 21)**

1 GGGATATTCT AGGACTTGCC TTGCAGAATC TGGTTGTTCT CCAAATGCCC  
51 TATGGAAGTG GAGAGCAGAA TGCTGCCCTA CTAGCATCTG ATACTTATGT  
101 TCTGGACTAT CTGAAATCTA CTGAGCAACT GACAGAGGAA GTTCAATCTA  
151 AGGCTTTCTT TCTCTTATCT AATG

10

**SEQ ID NO: 42 (INSP087 protein sequence exon 21)**

1 DILGLALQNL VVLQMPYGSQ EQNAALLASD TYVLDYLKST EQLTEEVQSK  
51 AFFLLSNG

**15 SEQ ID NO: 43 (INSP087 nucleotide sequence exon 22)**

1 GTTATCAAAG GCAATTATCT TTCAAAACT CTGATGGTTC CTATAGTGTG  
51 TTTTGGCAGC AGAGTCAGAA AGGAAGCATA TG

**SEQ ID NO: 44 (INSP087 protein sequence exon 22)**

20 1 YQRQLSEKNS DGSYSVFWQQ SQKGSIC

**SEQ ID NO: 45 (INSP087 nucleotide sequence exon 23)**

1 TGCTCTTACT TTTAAGACAT TGGAGAGAAT GAAAAAATAT GTATTCATTG  
51 ATGAAAATGT TCAAAAACAG ACCTTAATCT GGCTTTCAAG CCAACAGAAA  
25 101 ACAAGCGGCT GCTTTAAGAA TGATGGCCAG CTTTTCACC ACGCCTGGGA  
151 G

**SEQ ID NO: 46 (INSP087 protein sequence exon 23)**

1 ALTFKTLERM KKYVFIDENV QKQTLIWLSS QQKTSKCFKN DGQLFNHAW  
30

**SEQ ID NO: 47 (INSP087 nucleotide sequence exon 24)**

1 GGTGGAGATG AAGAGGACAT TTCACTCACT GCGTATGTTG TTGGGATGTT  
51 CTTTGAAGCT GGGCTCAATT TCACT

**SEQ ID NO: 48 (INSP087 protein sequence exon 24)**

1 GGDEEDISLT AYVVGMMFFEA GLNFT

**5 SEQ ID NO: 49 (INSP087 nucleotide sequence exon 25)**

1 TTTCCTGCTC TACGAAACGC ACTCTTTTGC CTTGAAGCGG CATTGGACAG  
51 TGGTGTCACT AATGGCTATA ATCATGCAAT TCTAGCTTAT GCTTTTGCCT  
101 TAGCTGGAAA AGAGAAGCAA GTGGAATCTT TACTCCAAAC CCTGGATCAA  
151 TCTGCCCCAA AACTAA

10

**SEQ ID NO: 50 (INSP087 protein sequence exon 25)**

1 FPALRNALFC LEAALDSGVT NGYNHAILAY AFALAGKEKQ VESLLQTLNQ  
51 SAPKLN

**15 SEQ ID NO: 51 (INSP087 nucleotide sequence exon 26)**

1 ATAATGTCAT CTA CTG GGA AGAGAAAGGA AACCCAAGAC AGAAGAATTT  
51 CCATCCTTTA TTCCCTGGGC ACCTTCTGCT CAGACTGAGA AGAGTTGCTA  
101 CGTGCTGTTG GCTGTCATTT CCCGAAAAT TCCTGACCTC ACCTATGCTA  
151 GTAAGATTGT GCAGTGGCTT GCCCAACGGA TGAATTCCCA TGGAGGCTTT

20 201 TCTTCCAACC AG

**SEQ ID NO: 52 (INSP087 protein sequence exon 26)**

1 NVIIYWERERK PKTEEFPSFI PWAPSAQTEK SCYVLLAVIS RKIPDLTYAS  
51 KIVQWLAQRM NSHGGFSSNQ

25

**SEQ ID NO: 53 (INSP087 nucleotide sequence exon 27)**

1 GAAACTGCAG TTTGTCTTCT TGCCATAACC CGCTACATAA CCC

**SEQ ID NO: 54 (INSP087 protein sequence exon 27)**

30 1 ETAVCLLAIT RYITQ

**SEQ ID NO: 55 (INSP087 nucleotide sequence exon 28)**

1 AGGGGCTCTT CTCTAAGGAT CAAAACACTG TCACCTTTAG CAGTGAAGGA  
51 TCCAGTGAGA TTTTCCAGGT TAACGGTCAT AACCGCCTAC TGGTCCAACG

101 TTCAGAAGTA ACACAGGCAC CTGGAGAATA CACAGTAGAT GTGGAAGGAC  
151 ACGGTTGTAC ATTTATCCAG

**SEQ ID NO: 56 (INSP087 protein sequence exon 28)**

5 1 GLFSKDQNTV TFSSEGSSEI FQVNGHNRL VQRSEVTQAP GEYTVDVEGH  
51 GCTFIQ

**SEQ ID NO: 57 (INSP087 nucleotide sequence exon 29)**

1 GCCACCCTTA AGTACAATGT TCTCCTACCT AAGAAGGCAT CTGGATTTTC  
10 51 TCTTTCCTTG GAAATAGTAA AGAACTACTC TTCGACTGCT TTTGACCTCA  
101 CAGTGACCCT CAA

**SEQ ID NO: 58 (INSP087 protein sequence exon 29)**

1 ATLKYNVLLP KKASGFSLSL EIVKNYSSTA FDLTVTLK  
15

**SEQ ID NO: 59 (INSP087 nucleotide sequence exon 30)**

1 ATACACTGGA ATTCGCAATA AATCCAGTAT GGTGGTTATA GATGTAAAAA  
51 TGCTATCAGG ATTTACTCCA ACCATGTCAT CCATTGAAGA G

20 **SEQ ID NO: 60 (INSP087 protein sequence exon 30)**

1 YTGIRNKSSM VVIDVKMLSG FTPTMSSIEE

**SEQ ID NO: 61 (INSP087 nucleotide sequence exon 31)**

1 CTTGAAAACA AGGGCCAAGT GATGAAGACT GAAGTCAAGA ATGACCATGT  
25 51 TCTTTTCTAC TTGAAAAT

**SEQ ID NO: 62 (INSP087 protein sequence exon 31)**

1 LENKGQVMKT EVKNDHVLFFY LEN

30 **SEQ ID NO: 63 (INSP087 nucleotide sequence exon 32)**

1 GGTTTTGGTC GAGCAGACAG TTTCCCTTTT TCTGTTGAGC AGAGCAACCT  
51 TGTGTTCAAC ATTCAGCCAG CCCAGCCAT GGTCTACGAT TATTACGAAA  
101 AAG

**SEQ ID NO: 64 (INSP087 protein sequence exon 32)**

1 GFGRADSFPF SVEQSNLVFN IQPAPAMVYD YYEKE

**5 SEQ ID NO: 65 (INSP087 nucleotide sequence exon 33)**

1 AAGAAATATGC CCTAGCTTTT TACAACATCG ACAGTAGTTC AGTTTCCCAG  
51 TGA

**SEQ ID NO: 66 (INSP087 protein sequence exon 33)**

10 1 EYALAFYNID SSSVSQ

**SEQ ID NO: 67 (INSP087 nucleotide sequence)**

1 GTTCCTCAGG CCAGATCTGA CCCACTGGCA TTTATTACAT TTTCTGCTAA  
51 AGGAGCCACT CTCAACCTGG AAGAGAGGAG ATCTGTGGCA ATCAGATCCA  
15 101 GAGAGAATGT GGTCTTCGTA CAGACTGATA AACCCACCTA CAAGCCTGGA  
151 CAGAAAGTTC ATATATTAAC ATTATTTTTA TTTTATTTTC AGTATCCAGT  
201 GATCACCCCTT CAGGATCCTC AAAACAATCG GATTTTTTCAA AGGCAAAATG  
251 TGA CTCTCTT CCGAAATATT ACCCAACTCT CGTTCCA ACT GATTT CAGAA  
301 CCAATGTTTG GAGATTACTG GATTGTTGTG AAAAGAACT CAAGGGAGAC  
20 351 AGTGACACAC CAATTTGCTG TTAAAAGATA TGTGCTGCCC AAGTTTGAAG  
401 TTACAGTCAA TGCACCACAA ACAGTAACTA TTTCAGATGA TGAATTCCAA  
451 GTGGATGTAT GTGCTAAGTA CAACTTTGGC CAACCTGTGC AAGGGGAAAC  
501 CCAAATCCGG GTGTGCAGAG AGTATTTTTTC TTCAAGCAAT TGTGAGAAAA  
551 ATGAAAATGA AATATGTGAG CAATTTATTG CACAGTTGGA AAATGGTTGT  
25 601 GTTTCTCAAA TTGTAAATAC AAAAGTCTTC CAACTCTACC GTTCGGGATT  
651 GTTCATGACA TTTCATGTCG CTGTAATTGT TACAGAATCT GGGACAGTTA  
701 TGCAGATCAG CGAGAAGACC TCAGTTTTTA TCACTCAATT GCTTGGA ACT  
751 GTAAACTTTG AGAACATGGA TACATTCTAT AGAAGAGGGA TTTCTTATTT  
801 TGGA ACTCTT AAATTTTCGG ATCCCAATAA TGTACCTATG GTGAACAAGT  
30 851 TGTTGCAACT GGAGCTCAAT GATGAATTTA TAGGAAATTA CACTACGGAT  
901 GAGAATGGCG AAGCTCAATT TTCCATTGAC ACTTCAGACA TATTTGATCC  
951 AGAGTTCAAC CTAAAAGCCA CATATGTTTCG ACCTGAGAGC TGCTATCTTC  
1001 CCAGCTGGTT GACGCCTCAG TACTTGGATG CTC ACTTCTT AGTCTCACGC  
1051 TTTTACTCCC GAACCAACAG CTTCTCTGAAG ATTGTTCCAG AACCAAAGCA

1101 GCTTGAATGT AATCAACAGA AGGTTGTTAC TGTGCATTAC TCCCTAAACA  
1151 GTGAAGCATA TGAGGATGAT TCCAATGTAA AGTTCTTCTA TTTGATGATG  
1201 GTAAAAGGAG CTATCTTACT CAGTGGACAA AAGGAAATCA GAAACAAAGC  
1251 CTGGAATGGA AACTTCTCGT TCCCAATCAG CATCAGTGCT GATCTGGCTC  
5 1301 CTGCAGCCGT CCTGTTTGTG TATACCCTTC ACCCCAGTGG GGAAATTGTG  
1351 GCTGACAGTG TCAGATTCCA GGTGACAAAG TGCTTTAAAC ACAAGGTTAA  
1401 CATAAAGTTC TCTAACGAGC AGGGCTTACC TGGTTCCAAT GCTAGTCTCT  
1451 GTCTTCAAGC GGCGCCTGTC TTATTCTGTG CCCTCAGGGC TGTGGATAGG  
1501 AATGTCCTTC TACTGAAATC TGAACAACAG CTGTCAGCTG AAAGTGTGTA  
10 1551 TAACATGGTT CCAAGTATAG AGCCGTATGG TTATTTCTAC CATGGCCTCA  
1601 ATCTTGATGA TGGCAAGGAA GACCCTTGCA TTCCTCAGAG GGATATGTTC  
1651 TACAATGGTT TATATTACAC ACCTGTAAGC AACTATGGGG ATGGAGATAT  
1701 CTATAATATT GTCAGGAACA TGGGTCTAAA AGTCTTTACC AATCTCCATT  
1751 ACCGAAAACC AGAAGTATGT GTGATGGAGA GAAGGCTGCC ACTCCCTAAG  
15 1801 CCGCTTTATC TGGAAACAGA AAATTATGGT CCAATGCGTA GTGTTCCGTC  
1851 TAGAATTGCA TGTAGAGGGG AGAATGCTGA CTATGTAGAA CAGGCTATAA  
1901 TTCAAACAGT AAGAACAAAC TTCCCAGAGA CATGGATGTG GGACCTCGTC  
1951 AGTGTGCGATT CCTCAGGCTC TGCCAATCTT TCGTTCCTCA TTCCTGATAC  
2001 GATAACCCAA TGGGAGGCAA GTGGCTTTTG TGTGAATGGT GACGTTGGAT  
20 2051 TTGGCATTTC CTCTACAACC ACTCTAGAAG TCTCCCAACC TTTCTTTATT  
2101 GAGATTGCCT CACCCTTTTC GGTGTTCAA AATGAACAAT TTGATTTGAT  
2151 TGTCAATGTC TTCAGCTACC GGAATACATG TGTAGAGATT TCTGTTCAAG  
2201 TGGAGGAGTC TCAGAATTAT GAAGCAAATA TTCATACCTT GAAAATCAAT  
2251 GGCAGTGAGG TTATTCAAGC TGGAGGGAGG AAAACAAACG TCTGGACTAT  
25 2301 TATACCTAAG AAATTGGGTA AAGTGAATAT CACTGTAGTT GCTGAGTCCA  
2351 AACAAAGCAG TGCTTGCCCA AATGAAGGAA TGGAGCAGCA AAAGCTAAAC  
2401 TGGAAAGACA CTGTGGTCCA AAGCTTCTTA GTAGAGCCTG AAGGTATTGA  
2451 AAAGGAAAGG ACCCAGAGTT TCCTTATCTG TACAGAAGGT GCCAAAGCCT  
2501 CCAAGCAGGG AGTTTTGGAC TTGCCAAACG ATGTAGTAGA AGGGTCAGCC  
30 2551 AGAGGCTTTT TCACTGTTGT GGGGGATATT CTAGGACTTG CCTTGCAGAA  
2601 TCTGGTTGTT CTCCAAATGC CCTATGGAAG TGGAGAGCAG AATGCTGCCC  
2651 TACTAGCATC TGATACTTAT GTTCTGGACT ATCTGAAATC TACTGAGCAA  
2701 CTGACAGAGG AAGTTCAATC TAAGGCTTTC TTTCTCTTAT CTAATGGTTA  
2751 TCAAAGGCAA TTATCTTTCA AAAACTCTGA TGGTTCCTAT AGTGTGTTTT



2801 GGCAGCAGAG TCAGAAAGGA AGCATATGTG CTCTTACTTT TAAGACATTG  
 2851 GAGAGAATGA AAAAATATGT ATTCATTGAT GAAAATGTTC AAAAACAGAC  
 2901 CTTAATCTGG CTTTCAAGCC AACAGAAAAC AAGCGGCTGC TTTAAGAATG  
 2951 ATGGCCAGCT TTTCAACCAC GCCTGGGAGG GTGGAGATGA AGAGGACATT  
 5 3001 TCACTCACTG CGTATGTTGT TGGGATGTTC TTTGAAGCTG GGCTCAATTT  
 3051 CACTTTTCCT GCTCTACGAA ACGCACTCTT TTGCCTTGAA GCGGCATTGG  
 3101 ACAGTGGTGT CACTAATGGC TATAATCATG CAATTCTAGC TTATGCTTTT  
 3151 GCCTTAGCTG GAAAAGAGAA GCAAGTGGAA TCTTTACTCC AAACCCTGGA  
 3201 TCAATCTGCC CCAAACTAA ATAATGTCAT CTACTGGGAA AGAGAAAGGA  
 10 3251 AACCCAAGAC AGAAGAATTT CCATCCTTTA TTCCCTGGGC ACCTTCTGCT  
 3301 CAGACTGAGA AGAGTTGCTA CGTGCTGTTG GCTGTCATTT CCCGGAATAA  
 3351 TCCTGACCTC ACCTATGCTA GTAAGATTGT GCAGTGGCTT GCCCAACGGA  
 3401 TGAATTCCCA TGGAGGCTTT TCTTCCAACC AGGAACTGC AGTTTGTCTT  
 3451 CTTGCCATAA CCCGCTACAT AACCCAGGGG CTCTTCTCTA AGGATCAAAA  
 15 3501 CACTGTCACC TTTAGCAGTG AAGGATCCAG TGAGATTTTC CAGGTTAACG  
 3551 GTCATAACCG CCTACTGGTC CAACGTTTCA AAGTAACACA GGCACCTGGA  
 3601 GAATACACAG TAGATGTGGA AGGACACGGT TGTACATTTA TCCAGGCCAC  
 3651 CCTTAAGTAC AATGTTCTCC TACCTAAGAA GGCATCTGGA TTTTCTCTTT  
 3701 CCTTGGAAT AGTAAAGAAC TACTCTTCGA CTGCTTTTGA CCTCACAGTG  
 20 3751 ACCCTCAAAT AACTGGAAT TCGCAATAAA TCCAGTATGG TGGTTATAGA  
 3801 TGTAATAATG CTATCAGGAT TTAATCCAAC CATGTCATCC ATTGAAGAGC  
 3851 TTGAAAACAA GGGCCAAGTG ATGAAGACTG AAGTCAAGAA TGACCATGTT  
 3901 CTTTTCTACT TGGAAAATGG TTTTGGTCGA GCAGACAGTT TCCCTTTTTC  
 3951 TGTTGAGCAG AGCAACCTTG TGTTCACAT TCAGCCAGCC CCAGCCATGG  
 25 4001 TCTACGATTA TTACGAAAAA GAAGAATATG CCCTAGCTTT TTACAACATC  
 4051 GACAGTAGTT CAGTTTCCCA GTGA

**SEQ ID NO: 68 (INSP087 protein sequence)**

1 VPQARSDPLA FIFSAKGAT LNLEERRSVA IRSRENVVVFV QTDKPTYKPG  
 30 51 QKVHILTLFL FLFQYPVITL QDPQNNRIFQ RQNVTSFRNI TQLSFQLISE  
 101 PMFGDYWIVV KRNSRETVTH QFAVKRYVLP KFEVTVNAPQ TVTISDDEFQ  
 151 VDVCAKYNFG QPVQGETQIR VCREYFSSSN CEKNENEICE QFIAQLENGC  
 201 VSQIVNTKVF QLYRSGLFMT FHVAVIVTES GTVMQISEKT SVFITQLLGT  
 251 VNFENMDTFY RRGISYFGTL KFSDPNNVPM VNKLLQLELN DEFIGNYTTD

301 ENGEAQFSID TSDIFDPEFN LKATYVRPES CYLPSWLTPQ YLDAHFLVSR  
351 FYSRTNSFLK IVPEPKQLEC NQOKVVTVHY SLNSEAYEDD SNVKFFYLMM  
401 VKGAILLSGQ KEIRNKAWNG NFSFPISISA DLAPAAVLV YTLHPSGEIV  
451 ADSVRFQVDK CFKHKVNIF SNEQGLPGSN ASLCLQAAPV LFCALRAVDR  
5 501 NVLLKSEQQ LSAESVYNMV PSIEPYGYFY HGLNLDDGKE DPCIPQRDMF  
551 YNGLYYTPVS NYGDGDIYNI VRNMGLKVFT NLHYRKPEVC VMERRLP LPK  
601 PLYLETENYG PMRSVPSRIA CRGENADYVE QAI IQTVRTN FPETWMWDLV  
651 SVDSSGSANL SFLIPDTITQ WEASGFCVNG DVGFGISSTT TLEVSQPF FFI  
701 EIASPFSVQ NEQFDLIVNV FSyrntCvei SVQVEESQNY EANIHTLKIN  
10 751 GSEVIQAGGR KTNVWTIIPK KLGKVNITVV AESKQSSACP NEGMEQQKLN  
801 WKDTVQSF L VEPEGIEKER TQSFLICTEG AKASKQGVLD LPNDVVEGSA  
851 RGGFTVVGDI LGLALQNLVV LQMPYGSSEQ NAALLASDTY VLDY LKSTEQ  
901 LTEEVSQKAF FLLSNGYQRQ LSFKNSDGSY SVFWQQSQKG SICALT FKT L  
951 ERMKKYVFID ENVQKQTLIW LSSQOKTSGC FKNDGQLFNH AWEGGDEEDI  
15 1001 SLTAYVVG MF FEAGLNFTFP ALRNALFCLE AALDSGVTNG YNHAILAYAF  
1051 ALAGKEKQVE SLLQTL DQSA PKLNNVIYWE RERKPKTEEF PSFIPWAPSA  
1101 QTEKSCYVLL AVISRKIPDL TYASKIVQWL AQRMNSHGGF SSNQETAVCL  
1151 LAITRYITQG LFSKDQNTVT FSSEGSSEIF QVNGHNRLLV QRSEVTQAPG  
1201 EYTV DVEGHG CTFIQATLKY NVLLPKKASG FSL SLEIVKN YSSTAFDLTV  
20 1251 TLKYTGIRNK SSMVVIDVKM LSGFTPTMSS IEELENKGQV MKTEVKNDHV  
1301 LFYLENGFGR ADSFPFSVEQ SNL VFNIQPA PAMVYDY EK EYALAFYNI  
1351 DSSSVSQ

**SEQ ID NO: 69 (INSP088 nucleotide sequence exon 1)**

25 1 CAGGTTTCCC TTGGCTTCTC CCCCTCCCAG CAGCTTCCAG GAGCAGAAGT  
51 GGAGCTGCAG CTGCAGGCAG CTCCCGGATC CCTGTGTGCG CTCCGGGCGG  
101 TGGATGAGAG TGTCTTACTG CTTAGGCCAG ACAGAGAGCT GAGCAACCGC  
151 TCT

**30 SEQ ID NO: 70 (INSP088 protein sequence exon 1)**

1 QVSLGFSPSQ QLPGA EVELQ LQAAPGSLCA LRAVDES VLL LRPDRELSNR  
51 S

**SEQ ID NO: 71 (INSP088 nucleotide sequence exon 2)**

1 GTCTATGGGA TGTTCATT CTGGTATGGT CACTACCCCT ATCAAGTGGC  
51 TGAGTATGAT CAGTGTCCAG TGTCTGGCCC ATGGGACTTT CCTCAGCCCC  
101 TCATTGACCC AATGCCCCAA GGGCATTCGA GCCAGCGTTC CATTATCTGG  
151 AGGCCCTCGT TCTCTGAAGG CACGGACCTT TTCAGCTTTT TCCGG

5

**SEQ ID NO: 72 (INSP088 protein sequence exon 2)**

1 VYGMFPFWYG HYPYQVAEYD QCPVSGPWDF PQPLIDPMPQ GHSSQRSIIW  
51 RPSFSEGTDL FSFR

10 **SEQ ID NO: 73 (INSP088 nucleotide sequence exon 3)**

1 GACGTGGGCC TGAAAATACT GTCCAATGCC AAAATCAAGA AGCCAGTAGA  
51 TTGCAGTCAC AGATCTCCAG AATACAGCAC TGCTATGGGT G

**SEQ ID NO: 74 (INSP088 protein sequence exon 3)**

15 1 DVGLKILSNA KIKKPVDCSH RSPEYSTAMG A

**SEQ ID NO: 75 (INSP088 nucleotide sequence exon 4)**

1 CAGGCGGTGG TCATCCAGAG GCTTTTGAGT CATCAACTCC TTTACATCAA  
51 GCAGAGGATT CTCAGGTCCG CCAGTACTTC CCAGAGACCT GGCTCTGGGA  
20 101 TCTGTTTCCT ATTGG

**SEQ ID NO: 76 (INSP088 protein sequence exon 4)**

1 GGGHPEAFES STPLHQAEDS QVRQYFPETW LWDLFPIG

25 **SEQ ID NO: 77 (INSP088 nucleotide sequence exon 5)**

1 TAACTCGGGG AAGGAGGCGG TCCACGTCAC AGTTCCTGAC GCCATCACCG  
51 AGTGGAAGGC GATGAGTTTC TGCACTTCCC AGTCAAGAGG CTTCGGGCTT  
101 TCACCCACTG TTGGACTAAC TGCTTTCAAG CCGTTCTTTG TTGACCTGAC  
151 TCTCCCTTAC TCAGTAGTCC GTGGGGAATC CTTTCGTCTT ACTGCCACCA  
30 201 TCTTCAATTA CCTAAAGGAT TGCATCAGG

**SEQ ID NO: 78 (INSP088 protein sequence exon 5)**

1 NSGKEAVHVT VPDAITEWKA MSFCTSQSRG FGLSPTVGLT AFKPFFVDLT  
51 LPYSVVRGES FRLTATIFNY LKDCIR

**SEQ ID NO: 79 (INSP088 nucleotide sequence exon 6)**

1 GTTCAGACTG ACCTGGCTAA ATCGCATGAG TACCAGCTAG AATCATGGGC  
51 AGATTCTCAG ACCTCCAGTT GTCTCTGTGC TGATGACGCA AAAACCCACC  
5 101 ACTGGAACAT CACAGCTGTC AAATTGG

**SEQ ID NO: 80 (INSP088 protein sequence exon 6)**

1 VQTDLAKSHE YQLESWADSQ TSSCLCADD A KTHHWNITAV KLG

**10 SEQ ID NO: 81 (INSP088 nucleotide sequence exon 7)**

1 GTCACATTAA CTTTACTATT AGTACAAAGA TTCTGGACAG CAATGAACCA  
51 TGTGGGGGCC AGAAGGGGTT TGTTCGCCAA AAGGGCCGAA GTGACACGCT  
101 CATCAAGCCA GTTCTCGTCA AA

**15 SEQ ID NO: 82 (INSP088 protein sequence exon 7)**

1 HINFTISTKI LDSNEPCGGQ KGFVPQKGRS DTLIKPVLVK

**SEQ ID NO: 83 (INSP088 nucleotide sequence exon 8)**

1 CCTGAGGGAG TCCTGGTGA GAAGACACAC AGCTCATTGC TGTGCCCAAA  
20 51 AG

**SEQ ID NO: 84 (INSP088 protein sequence exon 8)**

1 PEGVLVEKTH SLLCPKG

**25 SEQ ID NO: 85 (INSP088 nucleotide sequence exon 9)**

1 GAAAGGTGGC ATCTGAATCT GTCTCCCTGG AGCTCCAGT GGACATTGTT  
51 CCTGACTCGA CCAAGGCTTA TGTTACGGTT CTGG

**SEQ ID NO: 86 (INSP088 protein sequence exon 9)**

30 1 KVAESVSLE LPVDIVPDST KAYVTVLG

**SEQ ID NO: 87 (INSP088 nucleotide sequence exon 10)**

1 GAGACATTAT GGGCACAGCC CTGCAGAACC TGGATGGTCT GGTGCAGATG  
51 CCCAGTGGCT GTGGCGAGCA GAACATGGTC TTGTTTGCTC CCATCATCTA

101 TGTCTTGCAG TACCTGGAGA AGGCAGGGCT GCTGACGGAG GAGATCAGGT  
151 CTCGGGCAGT GGGTTTCCTG GAAATAG

**SEQ ID NO: 88 (INSP088 protein sequence exon 10)**

5 1 DIMGTALQNL DGLVQMPSGC GEQNMVLFAP IIVVLQYLEK AGLLTEEIRS  
51 RAVGFLEIG

**SEQ ID NO: 89 (INSP088 nucleotide sequence exon 11)**

1 GGTACCAGAA GGAGCTGATG TACAAACACA GCAATGGCTC ATACAGTGCC  
10 51 TTTGGGGAGC GAGATGGAAA TGGAAACACA TG

**SEQ ID NO: 90 (INSP088 protein sequence exon 11)**

1 YQKELMYKHS NGSYSAFGER DGNGNTW

15 **SEQ ID NO: 91 (INSP088 nucleotide sequence exon 12)**

1 GCTGACAGCG TTTGTCACAA AATGCTTTGG CCAAGCTCAG AAATTCATCT  
51 TCATTGATCC CAAGAACATC CAGGATGCTC TCAAGTGGAT GGCAGGAAAC  
101 CAGCTCCCCA GTGGCTGCTA TGCCAACGTG GGAAATCTCC TTCACACAGC  
151 TATGAAG  
20

**SEQ ID NO: 92 (INSP088 protein sequence exon 12)**

1 LTAEVTKCFG QAQKFIFIDP KNIQDALKWM AGNQLPSGCY ANVGNLLHTA  
51 MK

25 **SEQ ID NO: 93 (INSP088 nucleotide sequence exon 13)**

1 GGTGGTGTTG ATGATGAGGT CTCCTTGACT GCGTATGTCA CAGCTGCATT  
51 GCTGGAGATG GGAAAGGATG TAGAT

**SEQ ID NO: 94 (INSP088 protein sequence exon 13)**

30 1 GGVDDEVSLT AYVTAALLEM GKDVD

**SEQ ID NO: 95 (INSP088 nucleotide sequence exon 14)**

1 GACCCAATGG TGAGTCAGGG TCTACGGTGT CTCAAGAATT CGGCCACCTC  
51 CACGACCAAC CTCTACACAC AGGCCCTGTT GGCTTACATT TTCTCCCTGG

101 CTGGGGAAAT GGACATCAGA AACATTCTCC TTAAACAGTT AGATCAACAG  
151 GCTATCATCT CAG

**SEQ ID NO: 96 (INSP088 protein sequence exon 14)**

5 1 DPMVSQGLRC LKNSATSTTN LYTQALLAYI FSLAGEMDIR NILLKQLDQQ  
51 AIISG

**SEQ ID NO: 97 (INSP088 nucleotide sequence exon 15)**

1 GAGAATCCAT TTACTGGAGC CAGAAACCTA CTCCATCATC GAACGCCAGC  
10 51 CCTTGGTCTG AGCCTGCGGC TG TAGATGTG GAACTCACAG CATATGCATT  
101 GTTGGCCCAG CTTACCAAGC CCAGCCTGAC TCAAAGGAG ATAGCGAAGG  
151 CCACTAGCAT AGTGGCTTGG TTGGCCAAGC AACACAATGC ATATGGGGGC  
201 TTCTCTTCTA CTCAG

**15 SEQ ID NO: 98 (INSP088 protein sequence exon 15)**

1 ESIYWSQKPT PSSNASPWSE PAAVDVELTA YALLAQLTKP SLTQKEIAKA  
51 TSIVAWLAKQ HNAYGGFSST Q

**SEQ ID NO: 99 (INSP088 nucleotide sequence exon 16)**

20 1 GATACTGTAG TTGCTCTCCA AGCTCTTGCC AAATATGCCA CTACCGCCTA  
51 CATGCCATCT GAGGAGATCA ACCTGGTTGT AAAATCCACT GAGAATTTCC  
101 AGCGCACATT CAACATACAG TCAGTTAACA GATTGGTATT TCAGCAGGAT  
151 ACCCTGCCCA ATGTCCCTGG AATGTACACG TTGGAGGCCT CAGGCCAGGG  
201 CTGTGTCTAT GTGCAG

25

**SEQ ID NO: 100 (INSP088 protein sequence exon 16)**

1 DTVVALQALA KYATTAYMPS EEINLVVKST ENFORTFENIQ SVNRLVFQQD  
51 TLPNVPGMYT LEASGQGCY VQ

**30 SEQ ID NO: 101 (INSP088 nucleotide sequence exon 17)**

1 ACGGTGTTGA GATACAATAT TCTCCCTCCC ACAAATATGA AGACCTTTAG  
51 TCTTAGTGTG GAAATAGGAA AAGCTAGATG TGAGCAACCG ACTTCACCTC  
101 GATCCTTGAC TCTCACTATT CACACCAG

**SEQ ID NO: 102 (INSP088 protein sequence exon 17)**

1 TVLRYNILPP TNMKTFSLSV EIGKARCEQP TSPRSLTLTI HTS

**SEQ ID NO: 103 (INSP088 nucleotide sequence exon 18)**

5 1 TTATGTGGGG AGCCGTAGCT CTTCCAATAT GGCTATTGTG GAAGTGAAGA  
51 TGCTATCTGG GTTCAGTCCC ATGGAGGGCA CCAATCAGTT A

**SEQ ID NO: 104 (INSP088 protein sequence exon 18)**

1 YVGSRSSNM AIVEVKMLSG FSPMEGTNQL

10

**SEQ ID NO: 105 (INSP088 nucleotide sequence exon 19)**

1 CTTCTCCAGC AACCCCTGGT GAAGAAGGTT GAATTTGGAA CTGACACACT  
51 TAACATTTAC TTGGATGAG

**15 SEQ ID NO: 106 (INSP088 protein sequence exon 19)**

1 LLQQPLVKKV EFGTDTLNIY LDE

**SEQ ID NO: 107 (INSP088 nucleotide sequence exon 20)**

1 CTCATTAAGA AACTCAGAC TTACACCTTC ACCATCAGCC AAAGTGTGCT  
20 51 GGTCACCAAC TTGAAACCAG CAACCATCAA GGTCTATGAC TACTACCTAC  
101 CAG

**SEQ ID NO: 108 (INSP088 protein sequence exon 20)**

1 LIKNTQTYTF TISQSVLVTN LKPATIKVYD YYLPD

25

**SEQ ID NO: 109 (INSP088 nucleotide sequence exon 21)**

1 ATGAACAGGC AACAATTCAG TATTCTGATC CCTGTGAATG A

**SEQ ID NO: 110 (INSP088 protein sequence exon 21)**

30 1 EQATIQYSDP CE

**SEQ ID NO: 111 (INSP088 nucleotide sequence)**

1 CAGGTTTCCC TTGGCTTCTC CCCCTCCCAG CAGCTTCCAG GAGCAGAAGT  
51 GGAGCTGCAG CTGCAGGCAG CTCCCGGATC CCTGTGTGCG CTCCGGGCGG

101 TGGATGAGAG TGTCTTACTG CTTAGGCCAG ACAGAGAGCT GAGCAACCGC  
151 TCTGTCTATG GGATGTTTCC ATTCTGGTAT GGTCACCTACC CCTATCAAGT  
201 GGCTGAGTAT GATCAGTGTC CAGTGTCTGG CCCATGGGAC TTCCTCAGC  
251 CCCTCATTGA CCCAATGCCC CAAGGGCATT CGAGCCAGCG TTCCATTATC  
5 301 TGGAGGCCCT CGTTCTCTGA AGGCACGGAC CTTTTCAGCT TTTTCCGGGA  
351 CGTGGGCCTG AAAATACTGT CCAATGCCAA AATCAAGAAG CCAGTAGATT  
401 GCAGTCACAG ATCTCCAGAA TACAGCACTG CTATGGGTGC AGGCGGTGGT  
451 CATCCAGAGG CTTTTGAGTC ATCAACTCCT TTACATCAAG CAGAGGATTC  
501 TCAGGTCCGC CAGTACTTCC CAGAGACCTG GCTCTGGGAT CTGTTTCCTA  
10 551 TTGGTAACTC GGGGAAGGAG GCGGTCCACG TCACAGTTCC TGACGCCATC  
601 ACCGAGTGGA AGGCGATGAG TTTCTGCACT TCCCAGTCAA GAGGCTTCGG  
651 GCTTTCACCC ACTGTTGGAC TAACTGCTTT CAAGCCGTTT TTTGTTGACC  
701 TGACTCTCCC TTA CT CAGTA GTCCGTGGGG AATCCTTTTCG TCTTACTGCC  
751 ACCATCTTCA ATTACCTAAA GGATTGCATC AGGGTTCAGA CTGACCTGGC  
15 801 TAAATCGCAT GAGTACCAGC TAGAATCATG GGCAGATTCT CAGACCTCCA  
851 GTTGTCTCTG TGCTGATGAC GCAAAAACCC ACCACTGGAA CATCACAGCT  
901 GTCAAATTGG GTCACATTAA CTTTACTATT AGTACAAAGA TTCTGGACAG  
951 CAATGAACCA TGTGGGGGGCC AGAAGGGGTT TGTTCCCCAA AAGGGCCGAA  
1001 GTGACACGCT CATCAAGCCA GTTCTCGTCA AACCTGAGGG AGTCCTGGTG  
20 1051 GAGAAGACAC ACAGCTCATT GCTGTGCCCA AAAGGAAAGG TGGCATCTGA  
1101 ATCTGTCTCC CTGGAGCTCC CAGTGGACAT TGTTCCCTGAC TCGACCAAGG  
1151 CTTATGTTAC GGTCTCTGGGA GACATTATGG GCACAGCCCT GCAGAACCTG  
1201 GATGGTCTGG TGCAGATGCC CAGTGGCTGT GGCGAGCAGA ACATGGTCTT  
1251 GTTTGCTCCC ATCATCTATG TCTTGCACTA CCTGGAGAAG GCAGGGCTGC  
25 1301 TGACGGAGGA GATCAGGTCT CGGGCAGTGG GTTTCCTGGA AATAGGGTAC  
1351 CAGAAGGAGC TGATGTACAA ACACAGCAAT GGCTCATACA GTGCCTTTGG  
1401 GGAGCGAGAT GGAAATGGAA ACACATGGCT GACAGCGTTT GTCACAAAAT  
1451 GCTTTGGCCA AGCTCAGAAA TTCATCTTCA TTGATCCCCA GAACATCCAG  
1501 GATGCTCTCA AGTGGATGGC AGGAAACCAG CTCCCCAGTG GCTGCTATGC  
30 1551 CAACGTGGGA AATCTCCTTC ACACAGCTAT GAAGGGTGGT GTTGATGATG  
1601 AGGTCTCCTT GACTGCGTAT GTCACAGCTG CATTGCTGGA GATGGGAAAG  
1651 GATGTAGATG ACCCAATGGT GAGTCAGGGT CTACGGTGTC TCAAGAATTC  
1701 GGCCACCTCC ACGACCAACC TCTACACACA GGCCCTGTTG GCTTACATTT  
1751 TCTCCCTGGC TGGGGAAATG GACATCAGAA ACATTCTCCT TAAACAGTTA



1801 GATCAACAGG CTATCATCTC AGGAGAATCC ATTTACTGGA GCCAGAAACC  
1851 TACTCCATCA TCGAACGCCA GCCCTTGGTC TGAGCCTGCG GCTGTAGATG  
1901 TGGAACTCAC AGCATATGCA TTGTTGGCCC AGCTTACCAA GCCCAGCCTG  
1951 ACTCAAAAGG AGATAGCGAA GGCCACTAGC ATAGTGGCTT GGTGCGCCAA  
5 2001 GCAACACAAT GCATATGGGG GCTTCTCTTC TACTCAGGAT ACTGTAGTTG  
2051 CTCTCCAAGC TCTTGCCAAA TATGCCACTA CCGCCTACAT GCCATCTGAG  
2101 GAGATCAACC TGGTTGTAAA ATCCACTGAG AATTTCCAGC GCACATTCAA  
2151 CATAACAGTCA GTTAACAGAT TGGTATTTCA GCAGGATACC CTGCCCCAATG  
2201 TCCCTGGAAT GTACACGTTG GAGGCCTCAG GCCAGGGCTG TGTCTATGTG  
10 2251 CAGACGGTGT TGAGATACAA TATTCTCCCT CCCACAAATA TGAAGACCTT  
2301 TAGTCTTAGT GTGGAAATAG GAAAAGCTAG ATGTGAGCAA CCGACTTCAC  
2351 CTCGATCCTT GACTCTCACT ATTCACACCA GTTATGTGGG GAGCCGTAGC  
2401 TCTTCCAATA TGGCTATTGT GGAAGTGAAG ATGCTATCTG GGTTCAGTCC  
2451 CATGGAGGGC ACCAATCAGT TACTTCTCCA GCAACCCCTG GTGAAGAAGG  
15 2501 TTGAATTTGG AACTGACACA CTTAACATTT ACTTGGATGA GCTCATTAAG  
2551 AACACTCAGA CTTACACCTT CACCATCAGC CAAAGTGTGC TGGTCACCAA  
2601 CTTGAAACCA GCAACCATCA AGGTCTATGA CTACTACCTA CCAGATGAAC  
2651 AGGCAACAAT TCAGTATTCT GATCCCTGTG AATGA

20 **SEQ ID NO: 112 (INSP088 protein sequence)**

1 QVSLGFSPSQ QLPGALEVELQ LQAAPGSLCA LRAVDESVLL LRPDRELSNR  
51 SVYGMFPFWY GHYPYQVAEY DQCPVSGPWD FPQPLIDPMP QGHSSQRSII  
101 WRPSFSEGTDLFSFFRDVGL KILSNAKIKK PVDCSHRSPE YSTAMGAGGG  
151 HPEAFESSTP LHQAEDSQVR QYFPETWLWD LFPIGNSGKE AVHVTVPDAI  
25 201 TEWKAMSFCT SQSRGFGGLSP TVGLTAFKPF FVDLTLPLYSV VRGESFRLTA  
251 TIFNYLKDCI RVQTDLAKSH EYQLESWADS QTSSCLCADD AKTHHWNITA  
301 VKLGHINFTI STKILDSNEP CGGQKGFVPQ KGRSDTLIKP VLVKPEGVLV  
351 EKTHSSLLCP KGKVASESVS LELPVDIVPD STKAYVTVLG DIMGTALQNL  
401 DGLVQMPSGC GEQNMVLFAP IIYVLQYLEK AGLLTEEIRS RAVGFLEIGY  
30 451 QKELMYKHSN GSYSAFGERD GNGNTWLTAFTVKCFGQAQK FIFIDPKNIQ  
501 DALKWMAGNQ LPSGCIYANVG NLLHTAMKGG VDDEVSLTAY VTAALLEMGK  
551 DVDDPMVSQG LRCLKNSATS TTNLYTQALL AYIFSLAGEM DIRNILLKQL  
601 DQQAIIISGES IYWSQKPTPS SNASPWSEPA AVDVELTAYA LLAQLTKPSL  
651 TQKEIAKATS IVAWLAKQHN AYGGFSSTQD TVVALQALAK YATTAYMPSE

701 EINLVVKSTE NFQRTFNIQS VNRLVFQQDT LPNVPGMYTL EASGQGCYVYV  
751 QTVLRYNILP PTNMKTFSL S VEIGKARCEQ PTS PRSLT LT IHTSYVGSRS  
801 SSNMAIVEVK MLSGFSPMEG TNQLLLQQPL VKKVEFGTDT LNIYLD ELIK  
851 NTQTYTFTIS QSVLVTNLKP ATIKVYDY YL PDEQATIQYS DPCE

5

**SEQ ID NO:113 (INSP087 receptor binding domain protein sequence)**

MVVIDVKMLSGFTPTMSSIEELENKGQVMKTEVKNDHVL FYLENGFGRADSF PFSVEQSN  
LVFNIQPAPAMVYDYYEKEEYALAFYNIDSSSVSQ

10 **SEQ ID NO:114 (INSP087 receptor binding domain nucleotide sequence)**

ATGGTGGTTATAGATGTAAAAATGCTATCAGGATTTACTCCAACCATGTCATCCATTGAA  
GAGCTTGAAAACAAGGGCCAAGTGATGAAGACTGAAGTCAAGAATGACCATGTTCTTTTC  
TACTTGGAAAATGGTTTTGGTCGAGCAGACAGTTTCCCTTTTTCTGTTGAGCAGAGCAAC  
CTTGTGTTCAACATTCAGCCAGCCCCAGCCATGGTCTACGATTATTACGAAAAAGAAGAA

15 TATGCCCTAGCTTTTTACAACATCGACAGTAGTTCAGTTTCCCAG

**SEQ ID NO:115 (Cloned INSP087 receptor binding domain protein sequence)**

MVVIDVKMLSGFTPTMSSIEELENKGQVMKTEVKNDHVL FYLENGFGRADSF PFSVEQSN  
LVFNIQPAPAMVYDYYEKEEYALAFYNIDSSSVSE

20

**SEQ ID NO:116 (Cloned INSP087 receptor binding domain nucleotide sequence)**

ATGGTGGTTATAGATGTAAAAATGCTATCAGGATTTACTCCAACCATGTCATCCATTGAA  
GAGCTTGAAAACAAGGGCCAAGTGATGAAGACTGAAGTCAAGAATGACCATGTTCTTTTC  
TACTTGGAAAATGGTTTTGGTCGAGCAGACAGTTTCCCTTTTTCTGTTGAGCAGAGCAAC  
25 CTTGTGTTCAACATTCAGCCAGCCCCAGCCATGGTCTACGATTACTATGAAAAAGAAGAA  
TATGCCCTAGCTTTTTACAACATCGACAGTAGTTCAGTTTCCGAG

**SEQ ID NO:117 (INSP088 receptor binding domain protein sequence)**

NMAIVEVKMLSGFSPMEGTNQLLLQQPLVKKVEFGTDTLNIYLD ELIKNTQTYTFTISQS  
30 VLVTNLKPATIKVYDY YL PDEQATIQYSDPCE

**SEQ ID NO:118 (INSP088 receptor binding domain nucleotide sequence)**

AATATGGCTATTGTGGAAGTGAAGATGCTATCTGGGTTTCAGTCCCATGGAGGGCACCAAT  
CAGTTACTTCTCCAGCAACCCCTGGTGAAGAAGGTTGAATTTGGAAGTACACACTTAAC  
ATTTACTTGGATGAGCTCATTAAGAACTCAGACTTACACCTTCACCATCAGCCAAAGT  
5 GTGCTGGTCACCAACTTGAAACCAGCAACCATCAAGGTCTATGACTACTACCTACCAGAT  
GAACAGGCAACAATTCAGTATTCTGATCCCTGTGAA

**SEQ ID NO:119 (INSP088 macroglobulin domain protein sequence)**

10 QAEDSQVRQYFPETWLWDLFPIGNSGKEAVHVTVPDAITEWKAMSFCTSQSRGFGLSPTV  
GLTAFKPFVFDLTLPSVVRGESFRLTATIFNYLKDCIRVQTDLAKSHEYQLESWADSQT  
SSCLCADDAKTHHWNITAVKLGHINFITISKILDSNEPCGGQKGFVPQKGRSDTLIKPVL  
VKPEGVLVEKTHSSLLCPKGKVASESVSLELPVDIVPDSTKAYVTVLGDIMGTALQNLDG  
LVQMPSGCGEQNMVLFAPIIYVLQYLEKAGLLTEEIRSAVGFLEIGYQKELMYKHSNGS  
15 YSAFGERDGNNTWLTAFTKCFGQAQKFIFIDPKNIQDALKWMAGNQLPSGCYANVGNL  
LHTAMKGGVDDEVSLTAYVTAALLEMGKDVEDPMVSQGLRCLKNSATSTTNLYTQALLAY  
IFSLAGEMDIRNILLKQLDQQAIIISGESIYWSQKPTPSSNASPWSEPAVDVELTAYALL  
AQLTKPSLTQKEIAKATSIVAWLAKQHNAYGGFSSTQDTVVALQALAKYATTAYMPSEEI  
NLVVKSTENFQRTFNIQSVNRLVFQQDTLPNVPGMYTLEASGQGCYVYQTVLRYNILPPT  
20 NMKTFSLSVEIGKARCEQPTSPRSLTLTIHTSYVGSRSSNMAIVEVKMLSGFSPMEGTN  
QLLLQQPLVKKVEFGDTLNIYLDELIKNTQTYTFTISQSVLVTNLKPATIKVYDYLPD  
EQATIQYSDPCE

**SEQ ID NO:120 (INSP088 macroglobulin domain nucleotide sequence)**

25 CAAGCAGAGGATTCTCAGGTCCGCCAGTACTTCCCAGAGACCTGGCTCTGGGATCTGTTT  
CCTATTGGTAACTCGGGGAAGGAGGCGGTCCACGTCACAGTTCCTGACGCCATCACCGAG  
TGGAAGGCGATGAGTTTCTGCACTTCCCAGTCAAGAGGCTTCGGGCTTTCACCCACTGTT  
GGACTAACTGCTTTCAGCCGTTCTTTGTTGACCTGACTCTCCCTTACTCAGTAGTCCGT  
GGGGAATCCTTTCGTCTTACTGCCACCATCTTCAATTACCTAAAGGATTGCATCAGGGTT  
30 CAGACTGACCTGGCTAAATCGCATGAGTACCAGCTAGAATCATGGGCAGATTCTCAGACC  
TCCAGTTGTCTCTGTGCTGATGACGCAAAAACCCACCACTGGAACATCACAGCTGTCAA  
TTGGGTACATTAACCTTACTATTAGTACAAAGATTCTGGACAGCAATGAACCATGTGGG  
GGCCAGAAGGGGTTTGTTCCTCCAAAAGGGCCGAAGTGACACGCTCATCAAGCCAGTTCTC

GTCAAACCTGAGGGAGTCCTGGTGGAGAAGACACACAGCTCATTTGCTGTGCCCAAAGGA  
AAGGTGGCATCTGAATCTGTCTCCCTGGAGCTCCCAGTGGACATTGTTCCCTGACTCGACC  
AAGGCTTATGTTACGGTTCTGGGAGACATTATGGGCACAGCCCTGCAGAACCTGGATGGT  
CTGGTGCAGATGCCCAGTGGCTGTGGCGAGCAGAACATGGTCTTGTTTGTCTCCCATCATC  
5 TATGTCTTGCAGTACCTGGAGAAGGCAGGGCTGCTGACGGAGGAGATCAGGTCTCGGGCA  
GTGGGTTTCCCTGGAAATAGGGTACCAGAAGGAGCTGATGTACAAACACAGCAATGGCTCA  
TACAGTGCCTTTGGGGAGCGAGATGGAAATGGAAACACATGGCTGACAGCGTTTGTCA  
AAATGCTTTGGCCAAGCTCAGAAATTCATCTTCATTGATCCCAAGAACATCCAGGATGCT  
CTCAAGTGGATGGCAGGAAACCAGCTCCCCAGTGGCTGCTATGCCAACGTGGGAAATCTC  
10 CTTACACAGCTATGAAGGGTGGTGTGATGATGAGGTCTCCTTGACTGCGTATGTCACA  
GCTGCATTGCTGGAGATGGGAAAGGATGTAGATGACCCAATGGTGAGTCAGGGTCTACGG  
TGTCTCAAGAATTCGGCCACCTCCACGACCAACCTCTACACACAGGCCCTGTTGGCTTAC  
ATTTTCTCCCTGGCTGGGGAAATGGACATCAGAAACATTCTCCTTAAACAGTTAGATCAA  
CAGGCTATCATCTCAGGAGAATCCATTTACTGGAGCCAGAAACCTACTCCATCATCGAAC  
15 GCCAGCCCTTGGTCTGAGCCTGCGGCTGTAGATGTGGAACACAGCATATGCATTGTTG  
GCCCAGCTTACCAAGCCCAGCCTGACTCAAAGGAGATAGCGAAGGCCACTAGCATAGTG  
GCTTGGTTGGCCAAGCAACACAATGCATATGGGGGCTTCTCTTCTACTCAGGATACTGTA  
GTTGCTCTCCAAGCTCTTGCCAAATATGCCACTACCGCCTACATGCCATCTGAGGAGATC  
AACCTGGTTGTAAAATCCACTGAGAATTTCCAGCGCACATTC AACATACAGTCAGTTAAC  
20 AGATTGGTATTT CAGCAGGATACCCTGCCCAATGTCCCTGGAATGTACACGTTGGAGGCC  
TCAGGCCAGGGCTGTGTCTATGTGCAGACGGTGTGAGATACAATATTCTCCCTCCCACA  
AATATGAAGACCTTTAGTCTTAGTGTGGAAATAGGAAAAGCTAGATGTGAGCAACCGACT  
TCACCTCGATCCTTGACTCTCACTATTACACCAGTTATGTGGGGAGCCGTAGCTCTTCC  
AATATGGCTATTGTGGAAGTGAAGATGCTATCTGGGTT CAGTCCCATGGAGGGCACCAAT  
25 CAGTTACTTCTCCAGCAACCCCTGGTGAAGAAGGTTGAATTTGGAAC TGACACACTTAAC  
ATTTACTTGGATGAGCTCATTAAGAACACTCAGACTTACACCTTCACCATCAGCCAAAGT  
GTGCTGGTCACCAACTTGAAACCAGCAACCATCAAGGTCTATGACTACTACCTACCAGAT  
GAACAGGCAACAATTCAGTATTCTGATCCCTGTGAA

30 **SEQ ID NO:121 (Cloned fragment of INSP088 - protein sequence)**

GHPEAFESSTPLHQAEDSQVRQYFPETWLWDLFPIGNSGKEAVHVTVPDAITEWKAMSF  
TSQSRGFGLSPTVGLTAFKPFVVDLTLPYSVVRGESFRLTATIFNYLKDCIRVQTDLAKS  
HEYQLESWADSQTSSCLCADEAKTHHWNITAVKLGHINF TISTKILDSNEPCGGQKGFVP

QKGRSDTLIKPVLVKPEGVLVEKTHSSLLCPKGKVAESVSLELPVDIVPDSTKAYVTVL  
GDIMG TALQNLDGLVQMPSGCGEQNMVLFAPIIYVLQYLEKAGLLTEEIRSRAVG FLEIG  
YQKELMYKHSNGSYS AFGERDGNNGNTWLTAFVTKCFGQAQKFIFIDPKNIQDALKWMAGN  
QLPSG CYANVGNNLLHTAMKGGVDDEVSLTAYVTAALLEMGKDVDDPMVSQGLWCLKNSAT  
5 STTNLYTQALLAYIFSLAGEMDIRNILLKQLDQQAIIISGESIYWSQKPTPSSNASPWSEP  
AAVDVELTAYALLAQLTKPSLTQKEIAKATSIVAWLAKQRNAYGGFSSTQDTVVALQALA  
KYATTAYVPSEEINLVVKSTENFQRTFNIQSVNRLVFQQDTLPNVPGMYTLEASGQGC VY  
VQTVLRYNILPPTNMKTFSLSVEIGKARCEQPTSPRSLTLTIHTSYVGSRSSSNMAIVEV  
KMLSGFSPMEGTNQLLLQQPLVKKVEFGTDTLNIYLD ELIKNTQTYTFTISQSVLVTNLK  
10 PATIKVYDY YLPDEQATIQYSDPCE

**SEQ ID NO:122 (Cloned fragment of INSP088 – nucleotide sequence)**

GTGGTCATCCAGAGGCTTTTGAGTCATCAACTCCTTTACATCAAGCAGAGGATTCTCAGG  
TCCGCCAGTACTTCCCAGAGACCTGGCTCTGGGATCTGTTTCCTATTGGTAACTCGGGGA  
15 AGGAGGCGGTCCACGTCACAGTTCCTGACGCCATCACCGAGTGGAAGGCGATGAGTTTCT  
GCACTTCCCAGTCAAGAGGCTTCGGGCTTTCACCCACTGTTGGACTAACTGCTTTCAAGC  
CATTCTTTGTTGACCTGACTCTCCCTTACTCAGTAGTCCGTGGGGAATCCTTTCGTCTTA  
CTGCCACCATCTTCAATTACCTAAAGGATTGCATCAGGGTTCAGACTGACCTGGCTAAAT  
CGCATGAGTACCAGCTAGAATCATGGGCAGATTCTCAGACCTCCAGTTGTCTCTGTGCTG  
20 ATGAAGCAAAAACCCACCACTGGAACATCACAGCTGTCAAATTGGGTCACATTAAC TTTA  
CTATTAGTACAAAGATTCTGGACAGCAATGAACCATGTGGGGGCCAGAAGGGGTTTGTTC  
CCCAAAGGGCCGAAGTGACACGCTCATCAAGCCAGTTCTCGTCAAACCTGAGGGAGTCC  
TGGTGGAGAAGACACACAGCTCATTGCTGTGCCAAAAGGAAAGGTGGCATCTGAATCTG  
TCTCCCTGGAGCTCCCAGTGGACATTGTTCCCTGACTCGACCAAGGCTTATGTTACGGTTC  
25 TGGGAGACATTATGGGCACAGCCCTGCAGAACCTGGATGGTCTGGTGCAGATGCCCAGTG  
GCTGTGGCGAGCAGAACATGGTCTTGTTTGCTCCCATCATCTATGTCTTGCAGTACCTGG  
AGAAGGCAGGGCTGCTGACGGAGGAGATCAGGTCTCGGGCAGTGGGTTTCCTGGAAATAG  
GGTACCAGAAGGAGCTGATGTACAAACACAGCAATGGCTCATACAGTGCCTTTGGGGAGC  
GAGATGGAAATGGAAACACATGGCTGACAGCGTTTGT CACAAAATGCTTTGGCCAAGCTC  
30 AGAAATTCATCTTCATTGATCCCAAGAACATCCAGGATGCTCTCAAGTGGATGGCAGGAA  
ACCAGCTCCCCAGTGGCTGCTATGCCAACGTGGGAAATCTCCTTCACACAGCTATGAAGG  
GTGGTGTGATGATGAGGTCTCCTTGACTGCGTATGTCACAGCTGCATTGCTGGAGATGG  
GAAAGGATGTAGATGACCCAATGGTGAGTCAGGGTCTATGGTGTCTCAAGAATTCGGCCA  
CCTCCACGACCAACCTCTACACACAGGCCCTGTTGGCTTACATTTTCTCCCTGGCTGGGG

AAATGGACATCAGAAACATTCTCCTTAAACAGTTAGATCAACAGGCTATCATCTCAGGAG  
AATCCATTTACTGGAGCCAGAAACCTACTCCATCATCGAACGCCAGCCCTTGGTCTGAGC  
CTGCGGCTGTAGATGTGGAACACAGCATATGCATTGTTGGCCCAGCTTACCAAGCCCA  
GCCTGACTCAAAAGGAGATAGCGAAGGCCACTAGCATAGTGGCTTGGTTGGCCAAGCAAC  
5 GCAATGCATATGGGGGCTTCTCTTCTACTCAGGATACTGTAGTTGCTCTCCAAGCTCTTG  
CCAAATATGCCACTACCGCCTACGTGCCATCTGAGGAGATCAACCTGGTTGTAAAATCCA  
CTGAGAATTTCCAGCGCACATTCAACATACAGTCAGTTAACAGATTGGTATTTTCAGCAGG  
ATACCCTGCCCAATGTCCCTGGAATGTACACGTTGGAGGCCTCAGGCCAGGGCTGTGTCT  
ATGTGCAGACGGTGTTGAGATACAATATTCTCCCTCCCACAAATATGAAGACCTTTAGTC  
10 TTAGTGTGGAAATAGGAAAAGCTAGATGTGAGCAACCGACTTCACCTCGATCCTTGACTC  
TCACTATTACACCAGTTATGTGGGGAGCCGTAGCTCTTCCAATATGGCTATTGTGGAAG  
TGAAGATGCTATCTGGGTTTCAGTCCCATGGAGGGCACCAATCAGTTACTTCTCCAGCAAC  
CCCTGGTGAAGAAGGTTGAATTTGGAACGACACACTTAACATTTACTTGGATGAGCTCA  
TTAAGAACACTCAGACTTACACCTTCACCATCAGCCAAAGTGTGCTGGTCACCAACTTGA  
15 AACCAGCAACCATCAAGGTCTATGACTACTACCTACCAGATGAACAGGCAACAATTCAGT  
ATTCTGATCCCTGTGAA